

# A RECOVERY PLAN FOR THE CURTIS' PEARLY MUSSEL

---

*Epioblasma florentina curtisi*



A Recovery Plan for the Curtis' Pearly Mussel,  
Epioblasma florentina curtisi (Utterback 1915)

Prepared by

Alan C. Buchanan  
Missouri Department of Conservation  
1110 College Avenue  
Columbia, Missouri 65201

For

North Central Region  
U.S. Fish and Wildlife Service  
Twin Cities, Minnesota

Approved:

  
Regional Director, U.S. Fish and Wildlife Service

Date:

2-4-86

## Disclaimer

This is the completed Curtis' Pearly Mussel Recovery Plan. It has been approved by the U.S. Fish and Wildlife Service. It does not necessarily represent official positions or approvals of cooperating agencies and does not necessarily represent the views of all individuals, who played a role in preparing this plan. This plan is subject to modification as dictated by new findings, changes in species status, and completion of tasks described in the plan. Goals and objectives will be attained and funds expended contingent upon appropriations, priorities, and other constraints.

Acknowledgments should read as follows:

The Curtis' Pearly Mussel Recovery Plan, dated February 4, 1986, prepared by the U.S. Fish and Wildlife Service under contract with the Missouri Department of Conservation.

Literature Citations should read as follows:

U.S. Fish and Wildlife Service 1986. Curtis' Pearly Mussel Recovery Plan. U.S. Fish and Wildlife Service, Twin Cities, Minnesota. 92 pp.

Additional copies may be purchased from:

Fish and Wildlife Reference Service  
6011 Executive Boulevard  
Rockville, Maryland 20852  
301/770-3000  
or  
1-800-582-3421

## TABLE OF CONTENTS

	Page
I. Introduction .....	1
Description .....	1
Historical Distribution .....	3
Present Distribution .....	5
Ecology and Life History .....	8
Population Limiting Factors .....	10
Habitat .....	10
Hosts .....	11
Reproduction .....	11
Habitat Requirements .....	11
Endangered Species Status .....	12
Reason for Decline and Threats .....	12
II. Recovery .....	14
Recovery Plan Objective and Rationale .....	14
Step-down Outline .....	15
Narrative Outline .....	16
Literature Cited .....	22
III. Implementation Schedule .....	24
IV. Appendices .....	28

LIST OF FIGURES

Figure		Page
1.	Shell morphology of male (left) and female (right) <u>Epioblasma florentina curtisi</u> from Little Black River, Missouri .....	4
2.	Distribution of <u>Epioblasma florentina curtisi</u> (Utterback 1915) in Missouri .....	6
3.	Distribution of <u>Epioblasma florentina curtisi</u> (Utterback 1915) in Arkansas .....	7
4.	Typical naiad life cycle depicting various stages of development .....	9

## I. Introduction

The Curtis' Pearly Mussel, Epioblasma florentina curtisi, is on the verge of extinction. Although it has never been common or widespread, the range of E. f. curtisi has declined significantly since the early 1900s. Loss of habitat has been the primary cause of the decline of this species. E. f. curtisi requires shallow, flowing streams with stable substrates. Dam construction, sand and gravel dredging, and other stream disturbances have destroyed much of the habitat where this species originally occurred. Epioblasma florentina curtisi was listed as endangered on the United States List of Endangered and Threatened Wildlife and Plants (United States Department of the Interior 1976) on June 14, 1976, with an effective date of July 14, 1976.

Epioblasma florentina curtisi was first named Truncilla curtisi by Utterback (1915). Other synonyms include Dysnomia curtisi, and Plagiola turgidula. Clarke (1981) gives a brief explanation of the rationale for using Epioblasma rather than other generic names for this species.

### Description

The Curtis' Pearly Mussel was first described from specimens collected in the White River at Hollister, Missouri (Utterback 1915). His description was as follows:

**Nutritive Structures:** Branchial opening densely bordered with papillae; anal crenulated; supra-anal high, rather small, separated from anal by moderately short mantle connection, mantle border here spotted; inner laminae of inner gills entirely connected to visceral mass; palpi very small connected only by their anterior base which is remote from the

anterior attachment of pointed outer gill; color of soft parts grayish except the blackish mantle edge at siphonal openings.

**Reproductive Structures:** Marsupium very broad, rounded ventrad, kidney-shaped, ovisacs several, distinct occupying posterior half of outer gills and distended into beaded, unpigmented structures along ventral edge when gravid; conglutinates broken masses; glochidia unknown as only specimens gravid with early embryos have been obtained; inner mantle edge of female antero-ventrad to branchial opening drawn in toward the interior of shell forming a chamber."

**Shell Characters:** The shell is small, usually less than 1 1/2 inches (39 mm) in length. Males are slightly larger than females (mean length 31.5mm and 28.1 mm, respectively). The ratio of shell length to shell height to shell breadth is 6:4:3.

In the males, the shell is oval in shape, with the anterior end smoothly rounded, and the posterior end bluntly pointed and biangular. There is generally a slight but wide indentation on the posterior-ventral margin where the double, barely discernible posterior ridge joins the ventral margin (see Appendix A for a diagram illustrating shell terminology).

The female shell is smoothly rounded anteriorly and broadly rounded and inflated posteriorly. The posterior edge of the shell is serrated. The annual growth lines are deeply incised.

The shell of both sexes is yellowish brown to brown, usually light brown, sometimes with fine, evenly spaced rays over most of its length. The beaks are broad and low and the beak sculpture is typically eroded away. Utterback (1915) describes the beak sculpture as heavily ridged, tending towards double-looped.

The interior of the shell is white to whitish-blue, with the periostracum frequently showing through the nacre, particularly in the beaks and posterior half of the shell. Beak cavities are broad and relatively shallow. The pallial line, often indistinct when present, is one-third of the height of the shell above the ventral margin. The hinge line is broadly curved. Muscle scars are located against the hinge line. Anterior muscle scars are deeply impressed and touch the cardinal teeth, while posterior muscle scars are barely discernible. Two high, blade-like, short (one-third of the shell length) lateral teeth are found in the left valve, and one high blade-like lateral tooth is in the right valve. Lateral teeth are separated from the cardinal teeth by approximately half the laterals' length. Cardinal teeth are high, triangular, and divergent, one in the right valve and two in the left valve (Fig. 1).

#### Historical Distribution

Early in the 1900s, Epioblasma florentina curtisi occurred at scattered locations in the White and St. Francis River basins in southern Missouri and northern Arkansas (Figs. 2 & 3). In his "Naiadogeography of Missouri", Utterback (1917) listed this species as abundant in the White River Basin in southwestern Missouri and scarce in the Black River Basin in southeastern Missouri. Utterback (1915) reported this species from the White River near Hollister, Missouri. Johnson (1978) lists a collection of this species from the White River near Forsyth, Missouri in the early 1900s. The White River has been impounded at both Hollister and Forsyth, and E. f. curtisi no longer occurs in this stream. An old record (1916) of E. f. curtisi also exists from South Fork of the Spring River, a tributary to Black River in Arkansas (personal communication with Mr. Mark Gordon, University of Arkansas, Fayetteville, Arkansas).



Figure 1. Shell morphology of male (left) and female (right) *Epioblasma florentina curtisi* from Little Black River, Missouri (Photo by Nancy Lamb).

More recent historic records of E. f. curtisi exist for the Black River between Williamsville and Poplar Bluff in Wayne and Butler counties, Missouri (OSUM 13647 and 26509) (Oesch 1984; personal communication with Dr. David H. Stansbery, Ohio State University, Columbus, Ohio). These records are as recent as 1971.

#### Present Distribution

Since the mid-1970s E. f. curtisi has been found in the Castor River (Oesch 1984), Cane Creek, a tributary to the Black, and Little Black River (Fig. 2). Oesch (1984) found a subfossil shell of this species in Cane Creek in 1979; no fresh material was found. Living specimens and fresh shell of E. f. curtisi have been found in the Castor River at Zalma (OSUM 35918, 40480, 42223) between 1971 and 1978 (Oesch 1984; personal communication with Dr. David H. Stansbery, Ohio State Museum of Zoology, Columbus, Ohio; Bates and Dennis 1983). This species was found at 3 sites in 5 miles of the Upper Little Black River, a tributary to Current River, in 1979 during a survey funded by the Soil Conservation Service (Buchanan 1979). Bates and Dennis (1983) report finding 3 shells of an Epioblasma species from Black River at the mouth of Spring River in Arkansas. The specimens, however, are too different in size and other characteristics from Missouri specimens to be E. f. curtisi (personal communication with Mr. Mark Gordon, University of Arkansas, Fayetteville, Arkansas).

A survey was conducted during 1981 through 1983 to determine the distribution of the Curtis' Pearly Mussel in southern Missouri (Buchanan 1983). One hundred and forty-two sites were sampled on 26 streams between and including the James and Castor Rivers in southern Missouri. Fresh material of E. f. curtisi was found only in Little Black River. This species presently




Figure 2. Distribution of *Epioblasma florentina curtisi* (Utterback 1915) in Missouri.




Figure 3. Distribution of *Epioblasma florentina curtisi* (Utterback 1915) in Arkansas.

occurs only in 6.1 miles of the upper Little Black River and 7 miles or less of the Castor River upstream from the Headwater Diversion Channel (Fig. 2). It is known from only a total of six sites in these two streams.

### Ecology and Life History

Epioblasma florentina curtisi occurs in stream reaches which are transition areas between headwater and lowland stream reaches. It has been found in order 4 to 7 streams with gradients of 0.9 to 8.0 feet per mile, in stable substrates of sand and gravel to gravel, cobble and boulder, in riffles or runs (Buchanan 1983). Mean particle size of the substrate where this species is found, based on substrate analysis in the Little Black River, ranges from small gravel to cobble (Buchanan 1982). It is found in 2 to 30 inches of water in slow (less than 0.06 ft/sec. to 0.7 ft/sec. at the bottom) current. (Buchanan 1982; 1983). Other species of the genus Epioblasma also occur in shallow flowing water. Epioblasma walkeri (Wilson and Clark 1914) is found in coarse substrate in riffle areas of small to moderate sized rivers (Neves 1983). Ortmann (1919) found Epioblasma triquetra (Rafinesque 1829) and Epioblasma rangiana (Lea) in shallow water in strong current. He found E triquetra in a sand to coarse packed gravel substrate.

The Curtis' pearly mussel is assumed to have a life cycle similar to that of most other naiades<sup>1</sup> (Fig. 4). Males eject sperm directly into the water. The female holds eggs in her gills where they are fertilized by sperm taken in with water during siphoning. These eggs then develop into glochidia (the parasitic larval stage) in a modified portion of the gills called the marsupium. Fully developed glochidia are discharged into the water by female

---

<sup>1</sup> Naiades - freshwater bivalve molluscs with larvae (glochidia) which are parasitic on fish or other aquatic invertebrates.




Figure 4 Typical naiad life cycle depicting the various stages.

naiades. If a glochidium comes into contact with an appropriate fish host, it attaches to the fish's skin, fins, or gills and begins its parasitic stage.

Glochidia can neither swim nor crawl, and if a glochidium does not come into contact with a proper fish host, it falls to the bottom and dies. Each naiad species can successfully parasitize only certain species of fish. If a glochidium attaches to the wrong host, it will eventually be sloughed off and die. Some species have several fish hosts, while others may have only one. Most fish hosts are still unknown.

Glochidia live on the fish host for a week to several months, depending on the species, water temperature, and other factors. While attached, they receive nutrients from the fish. Eventually, glochidia develop into juvenile naiades and drop from the fish to begin life on the stream bottom.

Unfortunately the host for E. f. curtisi is not known.

Epioblasma florentina curtisi is bradytictic (eggs are fertilized in the fall and mature glochidia are released during early spring), based on ongoing research by the Missouri Department of Conservation. Females are extremely difficult to find except during March and April when they can be found lying on their sides on the surface of the substrate, releasing glochidia. During the remainder of the year they remain buried in the substrate. Other Epioblasma species also appear to be bradytictic (Ortmann 1919; Bogan and Parmalee 1983).

#### Population Limiting Factors

##### Habitat

Epioblasma florentina curtisi is primarily limited by habitat. It requires shallow flowing riffles and runs and relatively unpolluted water. Much of the original habitat in which this species occurred has been inundated by waters dammed for flood control, water supply, hydropower production, and

recreation (Buchanan 1983). Dam construction has impacted E. f. curtisi and other naiad species in several ways: elimination of flowing water habitat resulting in stagnation of bottom waters and accumulation of silt and organic matter; alteration of food supply; and possible elimination of fish hosts or restriction of their movement. Stream channelization and gravel dredging have reduced substrate variability and stability. Poor land management practices have increased siltation. Other forms of pollution have made many waters uninhabitable.

#### Hosts

E. f. curtisi may be limited by its host. Unfortunately, its host is not known. Research is currently being conducted by the Missouri Department of Conservation in cooperation with the United States Fish and Wildlife Service to determine the host and reproductive cycle of the Curtis' pearly mussel.

#### Reproduction

In streams such as the Castor River and Cane Creek, populations of E. f. curtisi may be so low that successful reproduction is rare. In 1978 it took four persons 22 man-hours to find a single living specimen in Castor River. Living specimens have not been found in Cane Creek.

#### Habitat Requirements

Like most naiades, E. f. curtisi requires flowing, relatively pristine silt-free waters. More specifically, this species appears to require shallow, stable riffles and runs in transitional stream reaches between headwater and lowland stream reaches. It has been found in order 4 to 7 streams with gradients of 0.9 to 8.0 feet per mile. Recently it has been found in 2 to 30

inches of water, in stable substrates of sand and gravel to gravel, cobble and boulder (Buchanan 1982; 1983).

#### Endangered Species Status

Epioblasma florentina curtisi was listed as endangered on the United States List of Endangered and Threatened Wildlife and Plants (United States Department of the Interior 1976) on June 14, 1976, with an effective date of July 14, 1976. This species remains extremely uncommon in the 15 miles or less of streams in which it occurs.

#### Reason for Decline and Threats

Habitat alteration has been and remains the principal threat to E. f. curtisi. This species requires shallow flowing water and a stable substrate. Consequently, stream impoundment and dredging have been the principal factors in the decline of E. f. curtisi. Impoundment caused the disappearance of E. f. curtisi from White River, impoundment and gravel dredging may have contributed to the decline of this species in Black River, and impoundment may pose a threat to this species in Little Black River (Buchanan 1983).

Populations of E. f. curtisi in White River have been eliminated by a series of impoundments constructed since the early 1900's. Lake Taneycomo, impounded in 1913, covered the sites where this species was found in the early 1900's (Utterback 1915; Johnson 1978). White River was further impounded by Bull Shoals Reservoir in 1952, Table Rock Lake in 1959, and Beaver Reservoir in 1966.

The disappearance of E. f. curtisi from Black River may be related to dam construction and gravel dredging (Buchanan 1983). Since closure of Clearwater Dam, flow characteristics in Black River downstream have changed drastically.

Hydrographic records at Leeper, 6.0 miles downstream from Clearwater Dam, reflect these changes (United States Geological Survey unpublished data). The dam has resulted in a drastic reduction in maximum annual flows, a reduction in mean daily flows in April, May, and June, and an increase in mean daily flows during July, August, and September. Clearwater Dam has caused much lower downstream flows than occurred before its construction. Very low flows may have exposed concentrations of naiades, causing desiccation, temperature stress, and increased susceptibility to predators. Changes in flow periodicity may have impacted reproduction through impacts on available fish hosts, water temperature, etc.

In the Little Black River, E. f. curtisi may be impacted by proposed reservoir construction 4 to 6 miles upstream. The impact of an impoundment on the aquatic organisms downstream is not easily predicted. Any effect which an impoundment has upon water temperature and chemical characteristics, types and amount of sediment, flow volume, duration and periodicity, and other factors will ultimately impact naiades. Changes in fish populations may also affect the chances of naiad reproduction. Changes in the microorganisms in the water may impact a naiad's food supply. Two impoundments proposed to be built upstream from populations of E. f. curtisi in Little Black River will cause changes in flow volume and periodicity, and may change the availability of nutrients and food, and cause an increase in scouring (Buchanan 1983).

Encroachment due to home construction and agricultural activities may impact E. f. curtisi in both the Little Black and Castor rivers. Construction of buildings on the floodplains of these two streams may result in bank destabilization and alteration of the channel to reduce flooding damage to those buildings. Agricultural activities in the floodplain can cause bank

destabilization and an increase in nutrient and sediment loads. Deforestation of the Little Black watershed may result in gravelling in of the river itself.

## II. Recovery

### Recovery Plan Objective and Rationale

The primary objective of this plan is to prevent the extinction of E. f. curtisi and restore its populations so that it can be removed from the Federal List of Endangered and Threatened Species. In order to do this, it is necessary to: 1) protect existing populations of this species and its habitat, 2) enhance existing populations and associated habitat, and 3) restore viable populations of E. f. curtisi to suitable habitat in its present and former range.

E. f. curtisi shall be considered recovered when: 1) viable<sup>2</sup> reproducing populations occur in four streams sufficiently separate so that a single catastrophic event would be unlikely to affect more than one population, 2) provisions have been made to prevent any future man-made changes (dam construction, channelization, in-stream dredging, etc.) that have the potential to adversely affect mussels in those streams (i.e. the riverways themselves are protected from significant changes due to industry, agriculture, or other forms of human encroachment and development), and 3) the four populations are protected by some form of riparian protected zone.

This plan is divided into two parts: Part A with a short-term goal to

---

<sup>2</sup> Viable population - A reproducing population that is large enough to maintain sufficient genetic variation to enable it to evolve and respond to natural habitat change. The number of individuals needed to meet this criterion will be determined as one of the recovery tasks (U.S. Fish and Wildlife Service 1984).

prevent extinction, and Part B with a long-term goal to provide for recovery of this species.

### Step-Down Outline

#### **A. Prevent Extinction**

Protect existing populations of E. f. curtisi and its habitat.

1. Further define the habitat needs of E. f. curtisi.
2. Review proposals for actions having the potential to adversely affect E. f. curtisi or its habitat.
3. Develop land protection plans.
4. Set aside riparian zones where activities which may impact E f. curtisi are prohibited.
5. Carry out other land protection actions.
6. Monitor habitat and populations in protected areas.

#### **B. Recovery**

1. Enhance existing populations through habitat and population management.
  - a. Improve status of habitat-limited populations.
    - (1) Evaluate habitat conditions.
    - (2) Develop methods to improve habitat.
    - (3) Implement habitat improvement methods.
  - b. Enhance populations declining due to low numbers or density of mature individuals.
    - (1) Identify populations with insufficient numbers or density.
      - (a) Determine population standing stocks.
      - (b) Determine if reproduction is taking place.

- (2) Determine how many individuals constitutes a viable population.
  - (3) Develop artificial culture/transplantation techniques.
  - (4) Enhance such populations with adults or artificially propagated juveniles.
- c. Evaluate habitat and population enhancement efforts.
2. Restore viable populations of E. f. curtisi to unoccupied former range.
    - a. Identify limiting factors in former E. f. curtisi habitat.
    - b. Eliminate limiting factors.
    - c. Introduce E. f. curtisi to former habitat.
    - d. Protect reintroduced populations.
    - e. Monitor sites where E. f. curtisi has been introduced.

### Narrative Outline

#### **A. Prevent Extinction**

Protect populations of E f. curtisi in the Little Black and Castor Rivers, Cane Creek, and other streams where this species may occur.

Prevent adverse alteration of existing habitat, via establishment of riparian protected zones, if necessary. The portions of Little Black and Castor Rivers where this species occurs are essential for its survival, and should be protected from adverse impacts resulting from development. Land protection planning should be initiated to determine the best means of preserving essential habitat areas.

1. Further define the habitat needs of E. f. curtisi so as to be able to better predict impacts of potential changes.
  - a. Continue to collect physical and chemical data at sites where the Curtis' pearly mussel occurs.

- b. Map stream reaches where this species occurs and mark locations where individuals occur in relation to depth, gravel bars, vegetation, flow, etc.
  - c. Compare plankton composition in streams to contents of Curtis' pearly mussel stomachs to determine food habits of this species.
  - d. Record seasonal flow, temperature and chemical characteristics where this species occurs and where it occurred previously. Document any changes in historic flow, temperature, and chemical regimes in streams from which this species has disappeared.
2. Review proposals for actions having the potential to adversely affect E. f. curtisi or its habitat, and require modifications which minimize or eliminate adverse effects.
  3. Develop land protection plans which identify the best methods of methods of protecting essential habitat.
  4. Set aside, through purchase, lease, or other means identified in the land protection plans, riparian zones where activities which may adversely impact E. f. curtisi are prohibited.
  5. Carry out other steps identified in land protection plans.
  6. Monitor habitat and populations in protected areas to determine if protective measures are effective.
    - a. Reexamine habitat characteristics every 3 to 5 years, and relate changes in population density to habitat changes.

## **B. Recovery**

1. Enhance existing populations through habitat and population management.
  - a. Improve status of habitat-limited populations. Determine which

marginal populations are limited by existing habitat, and improve the status of those populations.

- (1) Evaluate habitat conditions at all sites having marginal populations using the habitat needs data obtained from A1.
  - (2) Devise methods of reducing or eliminating those habitat factors which are suppressing E. f. curtisi population levels.
  - (3) Implement habitat improvement methods.
- b. Enhance status of density-limited populations. Where existing habitat is capable of supporting greater populations than now exist, but existing populations are so low that reproductive success is poor, enhance populations through introduction of additional individuals from viable populations (without harming viable populations) or by introducing juveniles produced through artificial culture. Propagation of E. f. curtisi through artificial culture should be attempted only after the techniques for artificial culture have been refined and there is a high likelihood that the juveniles will survive in the natural environment. Removal of gravid females and/or mature glochidia from viable populations for artificial culture should take place only from populations which will not be jeopardized by such removal.
- (1) Evaluate populations to identify those where recruitment is limited by low numbers or density of mature individuals.
    - (a) Determine population standing stocks by quadrat sampling or mark/recapture techniques. Repeat procedures every 3 to 5 years to determine population trends.
    - (b) Determine if reproduction is taking place by examining females for mature glochidia in early spring, and by

sieving the substrate for juveniles.

(2) Determine how many individuals constitutes a viable population.

(a) The present basis for determining how many individuals constitute a viable population is theoretical. Franklin (1980) postulates that at least 500 individuals are necessary for a population to maintain its genetic variability indefinitely. Smaller populations may lose their genetic ability to adapt to changing conditions. The following types of research will be required to determine the exact numbers necessary for the Curtis' pearly mussel to remain viable:

1. Chromosome mapping, electrophoresis, and other genetic research to determine how much genetic variation exists in each population, and between individuals in each population.
2. Once it is determined how many individuals are necessary to carry the present genetic variation, the population size required to perpetuate this variation can be determined based on mortality and recruitment rates, sex ratio, fecundity, etc.
3. Genetic research towards determining how many individuals constitutes a viable population can be accomplished only after Task B1b(3) is completed. Present populations are insufficient to permit removal of adequate numbers of individuals for genetic research.

(3) Develop artificial culture and transplantation techniques.

(a) The Tennessee Valley Authority (Isom & Hudson 1982; Hudson

& Isom 1984) is developing techniques for culturing glochidia and juveniles artificially. TVA has successfully raised the glochidia of several mussel species to juveniles in an artificial medium consisting of essential amino acids, vitamins, catfish blood, and antibiotics. They have also had limited success growing juveniles in river water piped into the laboratory.

(b) The above techniques can be employed to propagate E. f. curtisi only after they have been proven successful with closely related species, artificially propagated juveniles of those species are surviving and developing under natural conditions, and glochidia of E. f. curtisi have been raised to the juvenile stage under artificial conditions and shown to survive and grow under natural conditions.

(4) Enhance such populations with adults or juveniles produced through artificial propagation.

c. Periodically evaluate enhancement efforts to see if those efforts have been successful.

(1) Assess populations to determine if adult numbers are increasing.

(2) Monitor introduced individuals for survival.

(3) Examine introduced individuals for gravidity.

2. Restore viable populations of E. f. curtisi to present and former range.

a. Identify limiting factors in habitats where E. f. curtisi occurred previously but no longer occurs.

(1) Based on information collected in Section A1, identify differences between sites where E. f. curtisi occurs and where it no longer occurs.

- b. Eliminate limiting factors if possible.
- c. Introduce adults or juveniles to former habitat.
- d. Protect reintroduced populations using the methods identified in "A", as appropriate.
- e. Monitor sites every 3 to 5 years where E. f. curtisi has been reintroduced to evaluate success of reintroduction, using techniques from Section B1.
  - (1) Determine population standing stocks.
  - (2) Determine if reproduction is taking place.

### Literature Cited

- Bates, J. M. and S. D. Dennis. 1983. Mussel (naiad) survey--St. Francis, White, and Cache rivers, Arkansas and Missouri. U.S. Army Corps of Engineers, Memphis District. Contract No. DACW66-78-C-0147. 89 pp.
- Bogan, A. And P. Parmalee. 1983. Tennessee's rare mollusks, IN: Tennessee's rare wildlife, final report: TWRA, Tennessee Department of Conservation and Tennessee Heritage Program. June 1979. Univ. of Tenn., Knoxville. 360 pp.
- Buchanan, A. C. 1979. Mussels (naiades) of the Little Black River Basin in Missouri and Arkansas. United States Department of Agriculture, Soil Conservation Service, Columbia, Missouri. 69 pp.
- Buchanan, A. C. 1982. A study of Epioblasma florentina curtisi (Utterback 1915), the Curtis' Pearly Mussel, in the Upper Little Black River, Missouri. United States Department of Agriculture, Soil Conservation Service, Columbia, Missouri. 11 pp.
- Buchanan, A. C. 1983a. Distribution, abundance, status, and habitat of the Curtis' Pearly Mussel Epioblasma florentina curtisi in southern Missouri. United States Department of the Interior, Fish & Wildlife Service. Endangered Species Project SE-3-1. 10 pp.
- Buchanan, A. C. 1983b. An evaluation of the potential impacts of SCS structures B-9 and C-7 upon the Curtis' Pearly Mussel, Epioblasma florentina curtisi (Utterback 1915), in the Upper Little Black River, Missouri. Missouri Department of Conservation, 1110 College Avenue, Columbia, Missouri 65201. 18 pp.
- Clarke, A. H. 1981. Determination of the precise geographical areas occupied by four endangered species of freshwater mollusks. Final Report to U.S. Fish and Wildlife Service, Contract No. 14-16-003-81-019. 62 pp.
- Franklin, I. A. 1980. Evolutionary change in small populations. In Conservation Biology: An Evolutionary-Ecological Perspective M.E. Soule and B.A. Wilcox, eds. pp. 135-150. Sinauer Associates, Sutherland, Massachusetts.
- Hudson, R. G. and B. G. Isom. 1984. Rearing juveniles of the freshwater mussels (Unionidae) in a laboratory setting. The Nautilus 98(4):129-135.
- Isom, B. G. and R. G. Hudson. 1982. In vitro culture of parasitic freshwater mussel glochidia. The Nautilus 96 (4):147-151.
- Johnson, R. I. 1978. Systematics and zoogeography of Plagiola (= Dysnomia = Epicblasma), an almost extinct genus of freshwater mussels (Bivalvia: Unionidae) from Middle North America. Bulletin of the Museum of Comparative Zoology 148(6):239-321.

- Oesch, R. D. 1984. Missouri Naiades. A Guide to the Mussels of Missouri. Missouri Department of Conservation. 270 pp.
- Ortmann, A. E. 1919. A monograph of the naiades of Pennsylvania. Pt. 3. Systematic account of the genera and species. Mem. Carnegie Mus. 8 (1): 1-389, pl. 1-21.
- Rafinesque, C. S. 1820 Monographic des Coquilles Bivalves at Fluviatiles de la riviere Ohio. Ann. Sci. Phys. Bruxelles. 5:287-322.
- United States Department of the Interior, Fish and Wildlife Service. 1976. Federal Register. Part IV. Endangered and threatened wildlife and plants. Federal Register 41(208):47181-47198.
- United States Fish and Wildlife Service. 1983. Tan Riffle Shell Mussel Recovery Plan. U.S. Fish and Wildlife Service, Atlanta, Georgia. 48 pp.
- United States Fish and Wildlife Service. 1984. Rough Pigtoe Pearly Mussel Recovery Plan. U.S. Fish and Wildlife Service, Atlanta, Georgia. 51 pp.
- Utterback, W. I. 1915. The naiades of Missouri. American Midland Naturalist 4(3-8):200 pp.
- Utterback, W. I. 1917. Naiadogeography of Missouri. American Midland Naturalist 5(1):26-30.
- Wilson, C. B. and H. W. Clark. 1914. The mussels of the Cumberland River and its tributaries. U.S. Bureau of Fisheries Document No. 781. pp. 1-67.

III. IMPLEMENTATION SCHEDULE

GENERAL CATEGORIES FOR IMPLEMENTATION SCHEDULE (COLUMN 1)

Information Gathering - I or R (research)

1. Population status
2. Habitat status
3. Habitat requirements
4. Management techniques
5. Taxonomic studies
6. Demographic studies
7. Propagation
8. Migration
9. Predation
10. Competition
11. Disease
12. Environmental contaminant
13. Reintroduction
14. Other information

Management - M

1. Propagation
2. Reintroduction
3. Habitat maintenance and manipulation
4. Predator and competitor control
5. Depredation control
6. Disease control
7. Other management

Acquisition - A

1. Lease
2. Easement
3. Management agreement
4. Exchange
5. Withdrawal
6. Fee title
7. Other

Other - O

1. Information and education
2. Law enforcement
3. Regulations
4. Administration

Priorities in column four of the implementation schedule are assigned as follows:

1. Priority 1      An action that must be taken to prevent extinction or to prevent the species from declining irreversibly.
2. Priority 2      An action that must be taken to prevent a significant decline in species population/habitat quality, or some other significant negative impact short of extinction.
3. Priority 3      All other actions necessary to provide for full recovery of the species.

Abbreviations used in the implementation schedule, mainly in columns 7 and 8, are as follows:

ARGFC	Arkansas Game and Fish Commission
LE	Law Enforcement Division, U.S. Fish and Wildlife Service
MODOC	Missouri Department of Conservation
SE	Endangered Species Division, U.S. Fish and Wildlife Service
TBD	To be determined
TVA	Tennessee Valley Authority
USACE	U.S. Army Corps of Engineers
USFS	U.S. Forest Service, Mark Twain National Forest

RECOVERY PLAN IMPLEMENTATION SCHEDULE  
CURTIS' PEARLY MUSSEL

GEN. CAT.	PLAN TASK	TASK #	PRIORITY #	TASK DURATION	RESPONSIBLE AGENCY		FISCAL YEAR COSTS (EST.) (\$1,000's)			COMMENTS/NOTES	
					FWS	OTHER	FY 87	FY 88	FY 89		
R3	Define habitat needs	A1	1	Ongoing	3,8	SE, Research	MODOC	50	50	50	Low intensity research currently underway. Large scale study needed, estimated at 150K over 3 years
M3	Review habitat impacting proposals	A2	1	Ongoing	3	SE,HR	MODOC USACE USFS	1	1	1	
O4	Land protection planning	A3	1	2 years	3	SE	MODOC USFS	3	3		Includes planning for Little Black River, Castor River & Cane Creek
A?	Protect riparian zone	A4	1	Est. FY-90 completion	3	SE,RE	MODOC USACE USFS	TBD	TBD	TBD	Costs TBD by strategies developed in task A3
TBD	Other habitat protections	A5	1	TBD	3	SE,LE	MODOC USFS	TBD	TBD	TBD	" "
R1	Monitor protected populations	A6	1	Ongoing	3	SE	MODOC	5		5	Monitor populations every other year
R2	Evaluate habitat conditions	B1a(1)	2	2 years	3,8	SE Research	MODOC				Will begin following completion of task A1. Estimate \$25K total cost
R2	Methods to improve habitat	B1a(2)	2	TBD	3,8	SE Research	MODOC				Will commence after task B1a(1) completed
M3	Improve habitat	B1a(3)	2	TBD	3	SE	MODOC				Follows B1a(2)
R1 R6	Identify low-density populations	B1b(1)	2	2 years	3	SE	MODOC				Much necessary data should be available from tasks A6 and B1a(1). Further work will await completion of those tasks

RECOVERY PLAN IMPLEMENTATION SCHEDULE  
CURTIS' PEARLY MUSSEL

GEN. CAT.	PLAN TASK	TASK #	PRIORITY #	TASK DURATION	RESPONSIBLE AGENCY		FISCAL YEAR COSTS (EST.) (\$1,000's)			COMMENTS/NOTES
					REG. PROGRAM	OTHER	FY 87	FY 88	FY 89	
R6 R14	Determine how many individuals constitute a viable population	B1b(2)	2	TBD	3,8 SE, Research	TVA MODOC				To be initiated following completion of task B
R7 R13	Develop methods for artificial culturing and trans-plantation	B1b(3)	2	TBD	3,8 SE, Research	TVA				May be able to adapt methods currently being developed for other naiades by TVA
M1 M2	Augment populations	B1b(4)	2	TBD	3,8 SE, Research	MODOC USFS				To follow tasks B1b(1) and (3)
R1 R2	Monitor enhancement actions	B1c	2	TBD	3	MODOC				To follow tasks B1a(3) and B1b(4)
R2	Identify limiting factors in former range	B2a	3	3 years	3,4, 8 SE, Research	MODOC ARGFC				To follow task A1
M?	Eliminate limiting factors in former range	B2b	3	TBD	3,4 SE	MODOC ARGFC USACE				To follow task B2a
M2	Reintroductions into former range	B2c	3	TBD	3,4 SE	MODOC ARGFC USFS				To follow tasks B1b(3) and B2b
M3 A?	Protect reintroduction sites	B2d	3	Ongoing	3,4 SE,HR,LE	MODOC ARGFC				To follow task B2c
R1	Monitor reintroduced populations	B2e	3	Ongoing	3,4 SE	MODOC ARGFC				To follow task B2d


APPENDICES

APPENDIX A. Morphology of a naiad shell illustrating shell terminology

APPENDIX B. Three reports from completed research on the Curtis' Pearly Mussel


APPENDIX C. List of Reviewers

# Dorsal



# Ventral

# Dorsal



# Ventral

Appendix B.

The following reports have been included as appendices to give the reader more specific information on research already completed on the Curtis' Pearly

Mussel:

Buchanan, A. C. 1983. Distribution, abundance, status, and habitat of the Curtis' Pearly Mussel Epioblasma florentina curtisi in southern Missouri. United States Department of the Interior, Fish & Wildlife Service. Endangered Species Project SE-3-1. 10 pp.

Buchanan, A. C. 1982. A study of Epioblasma florentina curtisi (Utterback 1915), the Curtis' Pearly Mussel, in the Upper Little Black River, Missouri. United States Department of Agriculture, Soil Conservation Service, Columbia, Missouri. 11 pp.

Buchanan, A. C. 1983. An evaluation of the potential impacts of SCS Structures B-9 and C-7 upon the Curtis' Pearly Mussel, Epioblasma florentina curtisi (Utterback 1915) in the Upper Black River, Missouri. United States Department of Agriculture, Soil Conservation Service, Columbia, Missouri. 18 pp.

FINAL REPORT

Endangered Species Project SE-3-1  
Improving Status of Endangered Species in Missouri

DISTRIBUTION, ABUNDANCE, STATUS, AND HABITAT OF THE CURTIS' PEARLY  
MUSSEL EPIOBLASMA FLORENTINA CURTISI IN SOUTHERN MISSOURI

Missouri Department of Conservation

Larry R. Gale, Director

Charles A. Purkett, Jr., Assistant Director

Dan F. Dickneite, D-J Coordinator

James P. Fry, Chief, Division of Fisheries

Date Prepared:  
December 1, 1983

Prepared by:

Alan C. Suchanek  
Water Quality Research Biologist

Approved by:

James R. Hillery  
Fisheries Research Superintendent

## Abstract

One hundred and forty-six sites were sampled on twenty-six streams in southern Missouri during 1981 and 1982 to determine the distribution and status of Epioblasma florentina curtisi, a federally endangered species. Epioblasma florentina curtisi was not found in any new areas nor in areas of Black River and Cane Creek where it had been collected previously. No fresh specimens of E. f. curtisi have been found in Cane Creek or Black River since 1971. It occurs only in 6.1 miles of the Upper little Black River and 7 miles or less of the Castor River upstream from the Headwater Diversion Channel. These stream reaches should be considered critical habitat for E. f. curtisi. Most of the remaining E. f. curtisi occur in the Upper Little Black River. Management of this species should include periodic monitoring of populations, protection of critical habitat, searches for additional populations, reintroduction into areas from which it has disappeared, and introduction into other areas which appear suitable.

## Introduction

The Curtis Pearly Mussel, Epioblasma florentina curtisi, is one of those species which is so uncommon that it is on the verge of extinction. Although it has never been common or widespread, the range of Epioblasma florentina curtisi has declined significantly since the early 1900s. Loss of habitat has been the primary cause of the decline of this species. This species requires shallow, flowing streams with stable substrates. Dam construction, sand and gravel dredging, and other activities have destroyed much of the habitat where this species originally occurred.

The Curtis Pearly Mussel was first described from specimens collected in the White River at Hollister, Missouri (Utterback 1915). In his "Naiad-geography of Missouri" (1917) Utterback listed this species as "abundant" in the White River Basin in southwestern Missouri and "scarce" in the Black River Basin in southeastern, Missouri. Johnson (1978) lists collection of this species taken from the White River near Forsyth, Missouri in the early 1900s. The White River has been impounded at both of the above White River sites, and E. f. curtisi no longer occurs in this stream.

More recently E. f. curtisi has been found in the Black and Castor rivers (Oesch in press; personal communication with Dr. David H. Stansbery, Director, Museum of Zoology, Ohio State University, Columbus, Ohio), Cane Creek, a tributary to the Black River (Oesch in press), and in the Little Black River (Buchanan 1979). Living E. f. curtisi had only been collected in the Little Black and Castor rivers since the early 1970s.

This study was undertaken to gather information on the present distribution and status of E. f. curtisi in Missouri. The objectives of this study, conducted during April, 1981 through November, 1982 were: 1, to determine

the distribution and abundance of the Curtis Pearly Mussel; 2. to acquire information about its habitat; 3. to identify existing and potential threats to its survival; and 4. to develop recommendations for enhancing its status. This study was partially funded by the U. S. Department of the Interior, Fish and Wildlife Service.

#### Materials and Methods

Populations of Epioblasma florentina curtisi were assessed by field sampling in the Black, St. Francis, Castor, and White River basins in southern Missouri. Study sections included stream reaches where E. f. curtisi was found in the recent past and reaches believed likely to provide favorable habitat for this species, based on stream and basin characteristics such as area geology, topography, stream gradient, soil types, etc.

During April, 1981 through November, 1982, naiades were collected at 146 sites in 26 streams in southern Missouri (Fig. 1). Both stream reaches where E. f. curtisi had been found in the past and other reaches which appeared suitable for Epioblasma florentina curtisi were sampled. In most cases, a stream was sampled at 5-mile intervals in those reaches.

All habitat types were sampled at each site. Shells were collected by hand along shore, from muskrat and raccoon piles, and from the bottom of the stream. Live naiades were collected by hand while wading, and diving if necessary, until representatives of all species present were collected. The time spent collecting at each site ranged from 15 minutes to 7 1/2 hours; the average was 2 hours and 30 minutes.

The following information was recorded at each site: date, location, name(s) of collector(s), area covered by the site, bottom type(s) where




Figure 1. Sites sampled in southern Missouri.

living naiades were found, water temperature, water depth, collection techniques, the number of specimens of each species examined, and current velocity where living naiades were found. One or more specimens of all species except endangered species found at a site were retained as vouchers. Endangered species were photographed and returned unharmed to the habitat.

At each site, total hardness, pH, alkalinity, and dissolved oxygen were determined with a Hach kit, turbidity was measured with a Hach turbidimeter, and current velocity was measured with a Pygmy Current Meter. Current velocity was measured at mid-depth in water less than 6 inches deep and at surface and bottom in water greater than 6 inches deep.

## Results and Discussion

### Distribution and Abundance

Epioblasma florentina curtisi was not found in any new areas during this study. Also, no living specimens or fresh shell were found in areas of Black River and Cane Creek where it had been collected previously. No fresh material of E. f. curtisi has been found in Cane Creek or Black River since 1971. Based on this study, and a study on the upper Little Black River (Buchanan 1982), this species presently occurs only in 6.1 miles of the upper Little Black River and 7 miles or less of the Castor River upstream from the Headwater Diversion Channel (Fig. 2). It is known from a total of only six sites in these two streams. Following is a description of the distribution of E. f. curtisi in streams where it has been found both recently and historically.

### White River

No E. f. curtisi have been collected from White River since it was




Figure 2. Distribution of *Epioblasma florentina curtisi* (Utterback 1915) in Missouri.

impounded. Since E. f. curtisi is a riffle species, no sites were sampled on White River during this study. This species was not found in south-flowing tributaries sampled in southwestern Missouri during 1981-82, including James River, North Fork White River, and Swan, Bull, and Beaver creeks.

#### Black River

Epioblasma florentina curtisi has disappeared from Black River. It was first collected from Black River during the early 1900s (Utterback 1915). Utterback (1917) listed this species as "scarce" in the Black River Basin. During the 1960s E. f. curtisi was collected from Black River near Markham Spring, at Williamsville, and at Hendrickson (Oesch in press; personal communication with Dr. David H. Stansbery). Fresh shell of this species was collected from the Black River near Hendrickson in 1964 (OSUM 13647) and 1967 (OSUM 26509), and from Black River near Markham Spring in 1971 (OSUM 33633). No fresh material of this species has been found in Black River since 1971. No E. f. curtisi were found at 13 sites sampled between Clearwater Dam and Poplar Bluff during 1981 and 1982 (Fig. 1). The site near Hendrickson where it previously occurred has been dredged for sand and gravel. Old subfossil shell of this species was found in a muskrat midden near Williamsville but no fresh material was found.

The reason for the decline in E. f. curtisi populations in Black River is unknown. Changes in flow patterns in Black River since construction of Clearwater Dam in 1948 have probably contributed to the decline of E. f. curtisi in this stream. Hydrograph records at Leeper, 6.0 miles downstream from Clearwater Dam, reflect these changes (USGS). Between 1922 and 1948, the maximum annual flow was less than 10,000 cubic feet per second (cfs)

only 8 times and was less than 5,000 cfs only twice. Since construction of Clearwater Dam, the maximum annual flow at Leeper has never exceeded 10,000 cfs and has exceeded 5,000 cfs only twice. Periodicity of flows has been altered also. Mean daily flows at Leeper during April, May, and June were reduced by 357, 124, and 335 cfs, respectively, by Clearwater Dam. Mean daily flows during July, August, and September have been increased by 198, 160, and 126 cfs, respectively, by the dam. Clearwater Dam has also resulted in extremely low flow levels at Leeper. Between 1922 and 1948, annual minimum flows dropped below 200 cfs only 7 times, and the lowest flow was 133 cfs in August, 1934. Between 1949 and 1979 annual minimum flows dropped below 200 cfs 14 times and below 100 cfs twice. The lowest flows recorded were 70 cfs in November, 1959 and 62 cfs in September, 1966. The above flow changes could impact the time and bottom area available for fish, potential hosts for E. f. curtisi and other naiades, to spawn. Very low flows may expose concentrations of naiades, causing desiccation, temperature stress, and increased susceptibility to predators.

#### Cane Creek

A single subfossil shell of E. f. curtisi was collected in Cane Creek, a tributary to Black River, by Oesch (in press) in 1978. No other specimens of this species have been found in Cane Creek. None were found at three sites sampled between 21.5 miles and 29.0 miles upstream from the Arkansas border during 1980 and 1981. This stream changes rapidly from a headwater stream to a lowland stream and less than 5 miles of it provides potentially suitable habitat for E. f. curtisi. Naiad populations in Cane Creek are very limited. If E. f. curtisi still occurs in Cane Creek, it is present in very low numbers.

### Castor River

Utterback (1915, 1917) did not report finding E. f. curtisi in the Castor River. This species was first collected from the Castor River at Zalma during the 1960s by Frieda Schilling and Hessie Kemper (Oesch in press; personal communication with Dr. David H. Stansbery) (Fig. 2). The Ohio State University Museum (OSUM) has lots (OSUM 35918; OSUM 40480; OSUM 42223) containing a total of six specimens of E. f. curtisi collected at Zalma between 1973 and 1978. The Zalma site was last collected in 1978. At that time, E. f. curtisi occurred in very low numbers and only one specimen was found during 24 man-hours of effort. I felt no need to further disturb the site for the purposes of this study.

No E. f. curtisi were found at eight sites sampled in Castor River between 2.0 miles and 31.0 miles upstream from the Headwater Diversion Channel. Only about 6 miles, between 7.4 and 13.5 miles above the Headwater Diversion Channel, appear to provide suitable habitat for E. f. curtisi. Upstream from this 6-mile reach the stream becomes unstable, and downstream it becomes sluggish and lacks well-defined riffles. This species probably still occurs in Castor River in low numbers.

### Little Black River

The bulk of the remaining E. f. curtisi occurs in 6.1 miles of the upper Little Black River. Buchanan (1979) first found this species at three sites in 5 miles of the upper Little Black River during a survey of that stream funded by the Soil Conservation Service. During a subsequent study (Buchanan 1982) the range, habitat, and abundance of E. f. curtisi

in the upper Little Black River were further defined. Epioblasma florentina curtisi occurs between river miles 33.6 and 39.7 of this stream (Fig. 2).

Thirty living specimens of E. f. curtisi from the upper Little Black River, including 27 males and 3 females, were examined during 1979 through 1982, more specimens than have been found previously elsewhere. E. f. curtisi, however, is uncommon even in the Little Black River. Only 0.05 E. f. curtisi per square meter were found during quantitative sampling at three sites where this species is known to occur (Buchanan 1982). It comprised only 0.9 percent of the living naiades at those sites.

#### Habitat

Epioblasma florentina curtisi occurs in stream reaches which are transition areas between headwater and lowland stream reaches. It is found in order 4 to order 7 streams with gradients of 0.9 to 8.0 feet per mile, in stable substrates of sand and gravel to gravel, cobble and boulder, in riffles or runs. The mean particle size of the substrate where this species is found, based on substrate analysis in the Little Black River, ranges from small gravel to cobble (Buchanan 1982). It is found in 4 to 30 inches of water in slow (less than 0.02 m/sec. to 0.2 m/sec.) current.

#### Status

Epioblasma florentina curtisi is extremely uncommon in the less than 15 miles of streams where it occurs in southeastern Missouri. Only the upper Little Black River contains populations large enough to be effectively monitored and to serve as a source for reintroduction of E. f. curtisi into areas where it occurred previously. Loss of the Little Black River populations would probably result in extinction of the species. Epioblasma

florentina curtisi probably still occurs in Castor River in very low numbers. The Little Black River between river miles 33.6 and 39.7, and the Castor River between river miles 7.4 and 13.5 should be considered critical habitat for E. f. curtisi. Any impacts upon existing populations of this species may prevent its recovery.

#### Threats

Habitat alteration has been and remains the principal threat to E. f. curtisi. This species requires shallow flowing water and a stable substrate. Consequently, stream impoundment and dredging have been the principal factors in the decline of E. f. curtisi. Impoundment caused the disappearance of E. f. curtisi from White River and impoundment and gravel dredging contributed to the decline of this species in Black River.

In the Little Black River E. f. curtisi may be impacted by proposed reservoir construction 4 to 6 miles upstream. The impact of an impoundment on the aquatic organisms downstream is not easily predicted. Any effect which an impoundment has upon water temperature, the chemical parameters of water, the types and amount of sediment in the water, flow volume, duration and periodicity, and other factors will ultimately impact naiades. Changes in the fish present may affect the chances of a naiad successfully reproducing. Changes in the microorganisms in the water may impact a naiad's food supply.

Encroachment due to home construction and agricultural activities may impact E. f. curtisi in both the Little Black and Castor rivers. Construction of buildings on the floodplains of these two streams may result in bank destabilization and alteration of the channel to reduce flooding damage to those buildings. Agricultural activities in the floodplain can cause bank destabilization and an increase in nutrient and sediment loads.

#### Recommendations

1. The Little Black River between river miles 33.6 and 39.7 and the Castor River between 7.4 and 13.5 miles above the Headwater Diversion Channel should be listed as critical habitat for E. f. curtisi.
2. Populations of E. f. curtisi in the Little Black and Castor rivers should be monitored to detect changes which might affect the status of this species.
3. Portions of the habitat where E. f. curtisi occurs should be protected, through purchase or other means, to assure survival of existing populations.
4. Activities which might adversely impact E. f. curtisi in the Little Black and Castor rivers should be prevented.
5. Attempt to determine the reproductive cycle and host(s) of E. f. curtisi.
6. Continue to gather information on the habitat requirements for this species.
7. Identify stream reaches which may provide suitable habitat for E. f. curtisi.
8. Attempt to reestablish E. f. curtisi in habitats where it previously occurred. Care must be taken, however, to avoid harming existing populations by removal of too many individuals.

References Cited

- Buchanan, A. C. 1979. Mussels (naiades) of the Little Black River Basin in Missouri and Arkansas. United States Department of Agriculture, Soil Conservation Service, Columbia, Missouri. 69 pp.
- Buchanan, A. C. 1982. A study of Epioblasma florentina curtisi (Utterback 1915), the Curtis Pearly Mussel, in the Upper Little Black River, Missouri. United States Department of Agriculture, Soil Conservation Service, Columbia, Missouri . 11 pp.
- Johnson, R. I. 1978. Systematics and zoogeography of Plagiola (= Dysnomia = Epioblasma), an almost extinct genus of freshwater mussels (Bivalvia: Unionidae) from Middle North America. Bulletin of the Museum of Comparative Zoology 148(6):239-321.
- Oesch, R. D. In Press. The naiades of Missouri. Missouri Department of Conservation.
- United States Department of the Interior, Fish and Wildlife Service. 1976. Federal Register. Part IV. Endangered and threatened wildlife and plants. Federal Register 41(208):47181-47198.
- Utterback, W. I. 1915. The naiades of Missouri. American Midland Naturalist 4(3-8):200 pp.
- Utterback, W. I. 1917. Naiadogeography of Missouri. American Midland Naturalist 5(1):26-30.

A Study of Epioblasma florentina curtisi (Utterback 1915), the Curtis  
Pearly Mussel, in the Upper Little Black River, Missouri.

by Alan C. Buchanan

Missouri Department of Conservation  
Fish & Wildlife Research Center  
1110 College Avenue  
Columbia, Missouri

for the

U. S. Department of Agriculture  
Soil Conservation Service

February 1, 1982

## ABSTRACT

A study of the Curtis Pearly Mussel, Epioblasma florentina curtisi, in the Upper Little Black River, Missouri was conducted from May 1980 through November 1981. Twenty-one sites, including 11 sites on the Upper Little Black River, 2 sites on Flat Creek, 3 sites on Cane Creek, 1 site on Ten Mile Creek, and 4 sites on Castor River were examined. E. f. curtisi was not found in Flat or Ten Mile creeks or at any additional sites in Cane Creek or Castor River. The known range of E. f. curtisi was extended 1.2 miles in the Little Black River where it is now known from river mile 33.6 to river mile 39.7.

Comparison of naiad populations in the Upper Little Black River between river miles 33.6 and 43.5 revealed little difference in abundance, relative abundance, or species composition between 3 sites where E. f. curtisi is known to occur and 2 sites where this species does not occur. Seven species comprised 89.1% of the living naiades found in the 600 quadrats quantitatively sampled in the Upper Little Black River: Elliptio dilatata (29.3%), Pleurobema coccineum (23.7%), Lampsilis reeviana brevicula (15.6%), Ptychobranthus occidentalis (8.4%), Villosa lienosa lienosa (5.7%), Fusconaia flava (3.9%), and Toxolasma parvus (2.5%). Elliptio dilatata, P. coccineum, L. r. brevicula, P. occidentalis, and F. flava comprised over 75% of the living naiades found at both E. f. curtisi sites and non-E. f. curtisi sites. E. f. curtisi comprised only 0.6% of the naiades found. An average of 5.3 and 4.6 naiades/yd<sup>2</sup>, respectively, were found at E. f. curtisi sites and non-E. f. curtisi sites. There was no significant difference in water quality or substrate composition between sites where E. f. curtisi occurred and did not occur in the Upper Little Black River.

The 23 E. f. curtisi collected included 20 males and 3 females, ranging

from 0.6 to 1.5 inches long and from 4 to over 12 years old. Males were larger than females. They occurred in 4 inches to 30 inches of water, in slow (less than 0.2 yd/sec. to 0.26 yd/sec. at bottom) current, in a sand and gravel to a gravel, cobble and boulder substrate in or near riffles, in reaches of streams which serve as transition zones between headwaters and lowlands.

The populations of E. f. curtisi found in the Upper Little Black River are the best populations of this species known. The Little Black River between river miles 33.6 and 39.7 may be considered habitat essential to the survival of E. f. curtisi if no better populations of this species are found.

## INTRODUCTION

Epioblasma florentina curtisi, the Curtis Pearly Mussel, was first reported from the Upper Little Black River during a survey funded by the United States Department of Agriculture, Soil Conservation Service (Buchanan 1979). Although its presence was documented between river miles 33.6 and 38.5, its habitat requirements were not sufficiently described to evaluate the potential impacts of two proposed S.C.S. impoundments upon this species in the Upper Little Black River.

The primary objective of this study is to determine the distribution, abundance, habitat preference, and age structure of populations of E. f. curtisi in the Upper Little Black River and at selected sites in Cane Creek and the Castor River in Missouri. A secondary objective is to determine the reproductive success of E. f. curtisi in the above named streams. This study is being funded by the Columbia, Missouri office of the Soil Conservation Service to provide baseline data for evaluating the potential impacts of the two proposed S.C.S. impoundments (B-9 and C-7) upon the populations of E. f. curtisi in the Upper Little Black River.

## STUDY AREA

The primary study area, in the Little Black River Basin in southeastern Missouri (Fig. 1), is the Little Black River between river mile 43.5 (T24N, R3E, Section 4) in Ripley County and river mile 29.0 (T24N, R4E, Section 26) in Butler County (Fig. 2). The Little Black River in the primary study area is characterized by long, shallow, gravel and cobble-bottomed riffles interspersed with long, deep, silt and sand-bottomed pools. This portion of the Little Black River serves as a transition zone between its headwater and lowland reaches. Water willow (Justicia americana) is found in dense stands




Figure 1. Little Black River Basin, Missouri and Arkansas.




Figure 2. Sites sampled in the Upper Little Black River during 1980 and 1981.

along shore in riffle areas, and Spatterdock (Nuphar ozarkanum) and Pond week (Potamogeton sp.) are the most abundant plants in pools. The banks in most of the primary study reach are forested. Stream gradient ranges from 5.7 to 10.0 feet per mile.

The secondary study area is the Castor River between Highway 34 and the headwater diversion in Bollinger County, and Cane Creek between Highway PP and Highway 160 in Butler County (Fig. 3). These specific reaches of stream are transition zones between headwater and lowland streams habitats. Gradient in the study area of the Castor River ranges from 0.9 to 5.9 feet per mile. Cane Creek ranges in gradient from 2.5 to 5.9 feet per mile in the study reach. More specific water chemistry data for the above three stream reaches and Flat and Ten Mile creeks are listed in Table 6.

#### MATERIALS AND METHODS

Naiades were collected at 21 sites during May 1980 through November 1981, including 11 sites on the Little Black River, 4 sites on Castor River, 3 sites on Cane Creek, 2 sites on Flat Creek and 1 site on Ten Mile Creek (Fig. 3). Estimates of naiad density were made at 6 of the 21 sites (Fig. 3).

Initial sampling at 3 sites on the Castor River, 3 sites on Cane Creek, and 9 sites on the Little Black River was conducted to further delineate the distribution of E. f. curtisi in stream reaches from which it had previously been reported (Buchanan 1979) (including sites at approximately 1-mile intervals in the Little Black River between river miles 38.5 and 41.7 and between river miles 29.0 and 33.6). At these sites dead shells were collected by hand along shore, from muskrat and raccoon piles, and from the bottom of the stream. Live naiades were collected by hand while wading




Figure 3. Sites sampled in the Little Black and Castor rivers, and Flat, Cane, and Ten-Mile creeks during 1980 and 1981.

until we were confident that representatives of all species present had been collected.

Estimates of the density of naiades and Asiatic clams (number per square yard) were made at 5 sites on the Little Black River and 1 site on the Castor River. At each of these sites, a 50 (45m) to 110 (100m) yard stretch of stream which appeared to provide potential habitat for E. f. curtisi was marked off randomly (using a random numbers table) so that an average of one transect for every 3.3 yards (3m) of stream length was made. Along each transect, 1 to 7 quadrats (0.6 yd<sup>2</sup>) were selected (1 quadrat per every 5.5 yards (5m) of stream width). The substrate was sampled twice to a depth of 4 to 6 inches in each quadrat and all naiades and Asiatic Clams (Corbicula leana) were counted. Naiades and C. leana were counted in 600 quadrats on the Little Black River and in 92 quadrats in the Castor River.

Eleven to 20 randomly selected substrate samples were taken at the 5 sites on the Upper Little Black River where estimates of naiad densities were made. A 1-gallon can with one end removed was forced into the substrate as far as possible to obtain a substrate sample. These samples were frozen for later analysis.

Substrate samples were allowed to thaw and air-dry. Two randomly selected subsamples, each representing 1/8 of the original sample were taken for further analysis. The material was sieved to separate each subsample into nine different particle sizes (Table 3). Each size fraction was weighed, and its percent of the total weight of the subsample was calculated. The relative percent of each particle size was used to calculate mean Phi diameter, a measure of the mean particle size of the substrate (Inman 1952).

After sieving, all material less than 0.04 inches in diameter was combined from each subsample for determination of organic content. Each of these samples was dried at 100°C for 24 hours, weighed, ignited at 600°C for 2 hours and weighed again to determine percent loss at ignition.

Total hardness, pH, alkalinity, and dissolved oxygen were determined with a Hach kit, and turbidity was measured with a Hach turbidimeter at each site at which estimates of naiad density was made. These measurements were made quarterly during 1981. Current velocity was measured with a Pygmy Current Meter at locations where living E. f. curtisi were found. Current velocity was measured at mid-depth in water less than 6 inches deep, and at the surface and bottom in water greater than 6 inches deep.

#### RESULTS AND DISCUSSION

Epioblasma florentina curtisi was found 1.2 miles further upstream in the Little Black River during 1980 and 1981 than had been previously reported (Buchanan 1979). It was not, however, found any further downstream than previously reported. The Little Black River between river miles 30.4 and 33.6 appears to provide poor habitat for naiades and no naiad concentrations were found in that reach of the river. The known range of E. f. curtisi now extends from river mile 33.6 to river mile 39.7 (Fig. 2).

Twenty-three living specimens of E. f. curtisi, including 20 males and 3 females were found in the Upper Little Black River (Table 2). Males ranged from 0.6 to 1.5 inches long and from over 6 years old to over 12 years old. The 3 females were 0.9, 0.9, and 1.1 inches long and were all between 4 and 6 years old (Table 2). Twenty-one of the 23 specimens were found at river mile 37.9.

The composition of the substrate varied widely, both between sites and



Table 2. Age, sex, and size of the Epioblasma florentina curtisi found in Upper Little Black River during 1980 and 1981.

	River Mile	Sex	Length (inches)	Age (years)
1980	39.7	M	1.3	10+
		M	1.2	9
	37.9	M	1.2	10
		M	1.4	10
		M	1.5	12+
		F	1.1	5+
		M	0.6	6+
1981	37.9	M	1.2	9
		M	1.3	9+
		M	1.3	10
		M	1.2	8+
		M	1.3	8+
		M	1.2	9
		M	1.3	10+
		M	1.2	7+
		M	1.3	8
		M	1.3	9
		M	1.3	9
		M	1.3	12
		M	1.2	7+
		M	1.1	9
		F	0.9	4+
F	0.9	5		

at any one site (Table 3). There was no significant difference in the composition of the substrates between sites and no significant difference in the composition of the substrates where E. f. curtisi was found and where it was not. The mean particle size of the substrate samples ranged from coarse sand (0.02 to 0.04 inches in diameter) to cobble (1.25 to 2.5 inches in diameter). Organics comprised less than 0.6% of all substrate samples, and less than 0.4% of the substrate samples where E. f. curtisi was found.

Epioblasma florentina curtisi was found in 4 to 30 inches of water, in slow (less than 0.2 yd/sec. to 0.26 yd/sec. at bottom) current, in a sand and gravel to a gravel, cobble and boulder substrate. The mean particle size of the substrate where E. f. curtisi was found ranged from small gravel (0.04 to 0.16 inches in diameter) to pebble (0.62 to 1.25 inches in diameter).

An average of 5.1 naiades and 74.4 Corbicula leana per square yard was found in the Little Black River at the 5 sites where estimates of naiad density were made (Table 4, Fig. 2). Naiades were found in 63.8% of the quadrats sampled and occurred at all 5 sites. Corbicula leana was found in 72.0% of the quadrats sampled and occurred at 4 of 5 sites sampled; none were found at the most upstream site. Seventeen naiad species were found, including E. f. curtisi, and Cyprogenia aberti (Table 1.).

Seven species comprised 89.1% of the living naiades found in the 600 quadrats sampled in the Upper Little Black River (Table 4): Elliptio dilatata (29.3%), Pleurobema coccineum (23.7%), Lampsilis reeviana brevicula (15.6%), Ptychobranthus occidentalis (8.4%), Villosa lienosa lienosa (5.7%), Fusconaia flava (3.9%), and Toxolasma parvus (2.5%). Epioblasma florentina curtisi was found in 8 of the 600 quadrats sampled and comprised only 0.6% of the living naiades found. E. f. curtisi was found at only 1 of the 5 sites

Table 3. Particle size and organic composition of the substrate at five sites in the Upper Little Black River.

River mile	Transect	Depth (inches)	% to										% less than	% organic	Mean Phi diameter
			4.8 inches	2.50 inches	1.25 inches	0.62 inches	0.16 inches	0.04 inches	0.02 inches	0.01 inches	0.005 inches	0.01 inches			
36.7	3A	13	11.6	30.9	19.4	18.4	10.6	4.7	3.6	0.4	0.3	0.1	-3.45		
	4C	10	0.0	42.2	18.3	21.2	9.4	3.2	4.8	0.6	0.4	0.2	-3.68		
	10B	5	0.0	46.2	22.2	15.0	6.3	3.2	5.9	0.7	0.3	0.1	-3.90		
	12B	5	0.0	43.4	29.3	14.9	7.6	1.9	1.8	0.9	0.2	-	-3.98		
	27 Mid	7	20.3	28.5	10.6	20.4	10.4	4.6	4.0	0.9	0.2	-	-3.72		
	30E	7	0.0	22.1	14.0	24.7	9.8	8.0	18.4	1.4	0.4	0.2	-1.88		
	31D	7	0.0	23.6	15.5	26.3	13.6	8.4	11.8	0.5	0.2	0.1	-2.35		
	35D	6	0.0	22.6	31.4	24.2	8.9	5.3	7.1	0.4	0.1	0.1	-3.45		
	42E	5	0.0	24.3	20.6	21.7	13.8	7.2	9.9	1.9	0.6	0.2	-2.50		
	54 Left	12	25.9	19.4	18.4	18.4	7.3	3.8	5.0	1.4	0.3	-	-3.90		
	61A	14	0.0	3.9	27.9	27.6	11.4	8.3	16.2	3.7	0.8	-	-1.55		
	69A	17	12.6	24.0	17.8	18.3	9.8	7.1	9.0	0.8	0.5	0.2	-3.00		
	74C	14	11.2	10.4	20.6	35.2	7.2	4.1	9.4	1.3	0.4	0.3	-3.10		
	76D	9	0.0	12.4	27.6	32.1	10.8	5.7	9.2	1.9	0.3	0.1	-2.58*		
	77E	4	0.0	2.0	9.7	32.8	14.9	16.7	20.4	3.0	0.5	0.1	-1.10*		
	80E	2	0.0	0.0	7.3	31.3	15.7	19.9	24.2	1.1	0.4	0.3	-0.95		
	82B	17	0.0	27.6	27.2	22.7	9.7	5.6	5.1	1.6	0.5	-	-3.05		
93D	13	0.0	33.5	20.3	23.5	8.4	5.0	8.3	0.6	0.3	0.1	-3.10			
37.9	18C	22	0.0	53.5	19.5	16.9	4.6	1.4	2.7	1.3	0.2	0.1	-4.01*		
	2A	19	14.8	43.8	10.4	10.0	9.1	4.5	6.3	0.8	0.2	0.2	-3.85*		
	20B	17	0.0	65.8	13.4	11.1	3.4	1.5	3.9	0.6	0.2	0.1	-4.28		
	26A	18	0.0	0.0	0.0	24.0	34.6	22.2	15.9	2.7	0.5	-	-0.58		
	28C	23	9.6	46.7	12.3	19.4	6.9	1.2	2.9	0.6	0.3	0.2	-4.30*		
	38C	19	0.0	8.9	38.2	41.5	3.6	0.9	5.3	1.1	0.6	0.4	-3.62*		
	40A	17	0.0	25.7	4.6	28.7	18.4	11.3	10.0	0.8	0.5	0.3	-2.35*		
	41B	18	13.3	43.3	1.6	16.1	11.6	7.9	5.5	0.4	0.2	0.1	-3.42		
	45A	20	0.0	28.6	11.8	32.0	12.5	4.2	7.8	1.8	0.3	0.3	-2.88*		
	45B	22	6.7	23.6	6.7	21.4	15.2	11.9	11.4	2.6	0.5	0.3	-2.25*		
	45C	9	15.2	17.1	8.7	30.8	11.1	6.6	8.2	2.0	0.2	0.2	-3.05*		

Table 3 (continued). Particle size and organic composition of the substrate at five sites in the Upper Little Black River.

River mile	Transect	Depth (inches)	% to 4.8 inches	% to 2.50 inches	% to 1.25 inches	% to 0.62 inches	% to 0.16 inches	% to 0.04 inches	% to 0.02 inches	% to 0.01 inches	% less than 0.005 inches	% Phi	Mean diameter
38.5	-	13	0.0	22.6	8.9	21.0	17.4	9.7	12.0	7.2	1.0	0.2	-1.85*
39.7	3D	7	12.5	14.0	30.1	24.0	9.8	4.9	4.3	0.2	0.1	0.1	-3.75
	14E	11	0.0	42.9	15.9	16.6	9.2	7.3	7.2	0.4	0.4	0.3	-3.00
	19C	16	19.5	53.3	16.5	6.2	2.6	1.2	0.5	**	**	**	-5.20
	21F	13	0.0	18.0	14.7	31.2	8.1	10.0	16.4	1.0	0.5	0.4	-1.92
	23B	21	35.7	33.0	8.2	12.7	8.6	1.3	0.4	**	**	**	-4.70
	23D	11	59.2	21.6	8.5	5.8	2.0	1.3	1.4	**	**	**	-5.30
	28E	15	23.3	35.2	7.9	11.7	4.9	7.0	9.3	0.5	0.2	0.1	-3.10
	42D	21	0.0	49.1	9.1	23.9	11.1	4.4	2.0	2.0	0.2	0.2	-3.75
	52A	15	0.0	2.2	11.0	32.6	15.4	22.4	13.8	0.4	0.4	0.3	-1.30*
	55A	8	7.0	8.4	8.8	27.4	8.7	18.5	17.7	3.1	0.4	0.3	-1.80*
	58B	11	0.0	36.5	24.8	20.7	11.6	4.1	2.0	2.0	0.1	-	-3.70
	59C	18	18.9	24.3	24.5	17.5	9.4	4.0	0.1	1.2	0.1	0.1	-4.28
	61A	5	0.0	0.0	19.5	42.2	15.3	11.0	10.7	0.8	0.4	0.3	-1.80
	71B	9	42.1	21.6	14.6	12.9	6.7	1.5	0.5	0.5	**	-	-4.70
	74A	8	16.6	16.6	20.4	26.6	11.2	4.5	3.7	0.2	0.2	0.1	-3.90
	88B	13	0.0	21.1	19.0	25.5	23.7	7.2	3.1	3.1	0.3	0.1	-3.00
	89A	12	0.0	45.7	37.3	15.0	1.9	**	**	**	**	**	-4.60
	92C	8	18.7	20.2	20.3	18.0	12.1	5.5	4.6	4.6	0.4	0.2	-3.70
	95E	6	12.9	17.0	15.7	17.0	9.0	8.2	18.6	18.6	1.2	0.4	-2.85
	98E	6	0.0	40.0	14.3	23.7	5.8	3.6	10.8	10.8	1.3	0.3	-2.20
42.1	1F	15	0.0	19.4	6.0	28.9	12.7	13.6	17.3	1.3	0.9	0.4	-1.87
	3D	15	0.0	62.1	9.7	12.6	7.7	3.0	4.2	0.4	0.2	0.1	-3.90
	5A	9 1/2	0.0	0.0	23.9	21.2	11.2	21.1	20.2	1.2	1.0	0.6	-1.35
	12B	9	0.0	0.0	20.1	28.6	18.4	13.3	18.0	1.2	0.4	0.3	-1.45
	15D	5	0.0	16.9	22.7	21.4	17.1	8.0	11.0	2.6	0.3	-	-2.20
	17E	4 1/2	0.0	11.0	21.0	18.5	8.4	7.0	30.5	2.9	0.7	0.3	-1.50
	19C	6	0.0	50.6	11.5	15.5	10.1	4.2	7.0	0.7	0.3	0.2	-3.25
	20D	7 1/2	0.0	46.0	11.8	16.6	8.1	5.2	10.1	0.7	0.2	0.2	-2.52
	30C	8	25.3	38.8	5.6	15.3	9.1	2.7	2.5	0.5	0.1	-	-4.30

Table 3 (continued). Particle size and organic composition of the substrate at five sites in the Upper Little Black River.

River mile	Transect (inches)	Depth (inches)	% to										% less than	% organic	Mean Phi diameter
			4.8 inches	2.50 inches	1.25 inches	0.62 inches	0.16 inches	0.04 inches	0.02 inches	0.01 inches	0.005 inches	0.005 inches			
43.5	33B	4	11.9	10.2	12.8	22.4	18.1	11.3	11.3	11.3	1.9	0.2	-	-2.42	
	36C	6	36.5	8.2	10.5	23.6	11.0	4.4	5.2	0.4	0.2	0.1	-3.90		
	40D	6	0.0	24.9	3.6	29.0	16.5	9.4	15.3	0.9	0.4	0.2	-2.08		
	46E	5	0.0	17.3	28.0	17.6	10.3	5.9	17.5	2.9	0.4	-	-1.85		
	47B	7	0.0	11.0	21.0	18.5	8.4	7.0	30.5	2.9	0.7	0.2	-2.55		
	3C	14	0.0	8.8	2.0	0.7	6.9	20.0	47.1	11.9	2.6	-	+0.42		
	5E	9	0.0	46.2	12.2	20.6	13.0	3.4	3.5	0.6	0.5	0.2	-3.48		
8F	13	0.0	0.0	3.7	38.4	17.8	11.3	26.9	1.4	0.5	0.5	-0.72			
11D	7	0.0	34.6	14.1	25.9	15.5	3.7	4.4	1.3	0.4	-	-3.20			
17A	21	16.6	10.7	9.6	3.1	10.5	19.1	26.3	2.5	1.6	0.6	-2.18			
19E	21	22.8	40.3	8.3	11.9	5.4	3.4	6.4	1.0	0.6	0.2	-4.20			
20C	14	0.0	45.7	19.6	11.5	6.8	3.1	10.6	1.6	1.1	0.4	-2.28			
37E	14	0.0	39.9	14.9	11.4	8.6	7.5	14.7	1.7	1.1	0.6	-2.00			
40E	16	25.6	45.2	14.5	5.9	1.7	1.2	4.4	1.2	0.3	-	-5.05			
41A	13	11.1	15.7	10.1	32.9	15.1	7.9	6.0	0.9	0.2	-	-2.95			
42C	18	43.6	28.7	11.2	7.0	4.0	2.6	2.1	0.5	0.3	0.1	-4.45			
55E	6	0.0	28.0	19.0	25.5	13.9	5.0	7.3	0.8	0.5	0.2	-3.00			
57B	14	16.9	18.3	12.9	23.3	18.9	4.8	2.9	1.3	0.6	-	-3.55			
64C	15	39.6	24.9	9.3	11.5	6.0	4.1	3.8	0.6	0.2	-	-4.30			
73A	5	15.7	38.7	10.9	18.4	8.9	2.6	4.1	0.5	0.2	0.1	-4.10			
75D	18	33.8	31.6	11.3	11.1	4.5	2.3	4.3	0.8	0.2	-	-4.45			

\* *Epioblasma florentina curtisi* found in this substrate.

\*\* Less than 0.1%.

- Sample lost after sieving; no organic analysis possible.

Table 4. A comparison of naiades found in association with *Epioblasma florentina curtisi* in the Little Black River (quantitative samples only) during 1980 and 1981.

Species	Sites where <i>E.f. curtisi</i> found (three)			2 Sites where <i>E.f. curtisi</i> not found (two)			Total		
	Percent occurrence in quadrats	Number found	Relative abundance (%)	Percent occurrence in quadrats	Number found	Relative abundance (%)	Percent occurrence in quadrats	Number found	Relative abundance (%)
<i>Lasmigona costata</i>	2.2	10	0.8	2.9	8	1.1	2.5	18	* 0.9
<i>Tritogonia verrucosa</i>	2.2	8	0.6	0	0	0.0	1.3	8	* 0.4
<i>Amblema p. plicata</i>	6.1	34	2.8	0	0	0.0	3.7	34	1.8
<i>Fusconaia flava</i>	10.5	50	4.1	10.9	26	3.7	10.7	76	0.2
<i>Cyclonaias tuberculata</i>	4.7	21	1.7	6.7	20	2.8	5.5	41	0.1
<i>Pleurobema coccineum</i>	39.1	346	28.2	24.7	111	15.7	33.3	457	23.7
<i>Elliptio dilatata</i>	25.2	378	30.8	18.0	188	26.7	22.3	566	1.5
<i>Ptychobranchus occidentalis</i>	14.7	102	8.3	16.7	60	8.5	15.5	162	0.4
<i>Cyprogenia aberti</i>	4.2	23	1.9	2.5	7	1.0	3.5	30	0.1
<i>Actinonaias ligamentina carinata</i>	0.3	1	0.1	0	0	0.0	0.2	1	* *
<i>Toxolasma parvum</i>	3.6	26	2.1	5.8	23	3.3	4.5	49	0.1
<i>Villosa l. iris</i>	3.0	13	1.1	5.8	18	2.6	4.2	31	0.1
<i>Villosa l. lienosa</i>	8.3	91	7.4	5.4	19	2.7	7.2	110	0.3
<i>Lampsilis radiata luteola</i>	0.8	3	0.2	0.4	1	0.1	0.7	4	* 0.2
<i>Lampsilis ventricosa</i>	5.0	23	1.9	2.9	8	1.1	4.2	31	0.1
<i>Lampsilis reeviana brevicula</i>	16.3	86	7.0	51.0	216	30.6	30.2	302	0.8
<i>Epioblasma florentina curtisi</i>	2.2	11	0.9	0	0	0.0	1.3	11	* 0.6
Naiad totals	57.9	1,226	100.0	72.8	705	100.0	63.8	1,931	5.1
<i>Corbicula leana</i>	93.9	22,160	96.5	38.9	6,216	40.9	72.0	28,376	74.4
Area sampled	4,437.7 yd <sup>2</sup>			3,009.5 yd <sup>2</sup>			7,447.2 yd <sup>2</sup>		
Sample size	229.6 yd <sup>2</sup>			152.0 yd <sup>2</sup>			381.6 yd <sup>2</sup>		
Percent of area sampled	5.2%			5.1%			5.1%		
Length of stream sampled	300 yd			150 yd			450 yd		

\* = less than 0.1%

during quantitative sampling in 1980 and 1981, even though it had previously been found at 3 of these 5 sites (Buchanan 1979). This species comprised only 1.6% of the living naiades found at that site.

The mean density of naiades ranged from 1.9 to 21.7 naiades per square yard at river miles 36.7 and 37.9, respectively. There was little difference, however, between the mean density of naiades at the 3 sites at which E. f. curtisi occurs and the 2 sites where it does not occur (5.3/yd<sup>2</sup> and 4.6/yd<sup>2</sup>, respectively). E. f. curtisi was most abundant at the site which had the greatest concentration of all naiades.

There were few differences in species composition and relative abundance between the E. f. curtisi sites and the non-E. f. curtisi sites. Five species, E. dilatata, P. coccineum, P. occidentalis, L. r. brevicula, and F. flava comprised 78.4% of the living naiades found at the 3 E. f. curtisi sites and 85.2% of the living naiades found at the 2 non-E. f. curtisi sites (Table 4). L. r. brevicula, typically a headwater species, was the most abundant species at river mile 43.5 (Table 5). It comprised a progressively smaller portion of the naiad population, however, further downstream. Elliptio dilatata and P. coccineum were the most abundant species at the 3 E. f. curtisi sites and were second and third in abundance, respectively, at the 2 non-E. f. curtisi sites (Table 4).

Measures of water quality were similar at all sites in the Little Black River (Tables 6 and 7). Dissolved oxygen ranged from 5.5 to 13.2 mg/l, alkalinity from 0 to 137 mg/l, total hardness from 86 to 205 mg/l, pH from 7.0 to 8.5, and specific conductance from 110 to 165  $\mu$ mhos/cm<sup>2</sup>. No distinctive trends were evident between upstream and downstream portions of the primary study area.

Table 5. Numbers and relative abundance of naiad species found at each site sampled quantitatively in the Upper Little Black River during 1980 and 1981.

Species	River Mile 36.7			River Mile 37.9			River Mile 39.7		
	Percent occurrence in quadrats	Number found	Relative abundance (%)	Percent occurrence in quadrats	Number found	Relative abundance (%)	Percent occurrence in quadrats	Number found	Relative abundance (%)
<i>Lamigona costata</i>				12.2	8	0.3	1.2	2	0.6
<i>Tritogonia verrucosa</i>				14.3	7	0.2	1.0	1	0.3
<i>Amblema p. plicata</i>				42.9	33	1.1	4.9	*	
<i>Fusconaiia flava</i>	0.6	1	0.5	46.9	34	1.1	5.0	10	2.8
<i>Cyclonaiias tuberculata</i>	3.8	6	3.1	10.2	6	0.2	0.9	11	3.1
<i>Pleurobema coccineum</i>	2.5	4	2.1	77.6	157	5.0	23.3	75	20.9
<i>Elliptio dilatata</i>	41.5	114	59.1	93.9	307	9.8	45.5	42	11.7
<i>Psychobranchus occidentalis</i>	14.5	29	15.0	26.5	19	0.6	2.8	68	19.0
<i>Cyprogenia aberti</i>	8.8	15	7.8	8.2	4	0.1	0.6	18	5.0
<i>Actinonaiias ligamentina carinata</i>	0.6	1	0.5	2.0	1	*	0.1	20	5.6
<i>Toxolasma parvus</i>				10.2	6	0.2	0.9	1	0.3
<i>Villosa l. iris</i>	3.1	6	3.1	10.2	6	0.2	0.9	68	19.0
<i>Villosa l. lienosa</i>	3.8	6	3.1	16.3	17	0.5	2.5	0.7	
<i>Lampsilis radiata luteola</i>				6.1	3	0.1	0.4	8	2.2
<i>Lampsilis ventricosa</i>				26.5	15	0.5	2.2	34	9.5
<i>Lampsilis reeviana brevicula</i>				51.0	41	1.3	6.1		
<i>Epioblasma florentina curtisi</i>	6.3	11	5.7	16.3	11	0.4	1.6		
<b>Naiad totals</b>	<b>56.6</b>	<b>193</b>	<b>1.9 ± 0.3 c.i.</b>	<b>100.0</b>	<b>675</b>	<b>21.7 ± 4.1 c.i.</b>		<b>358</b>	<b>3.7 ± 1.2 c.i.</b>
<b>Corbicula leana</b>		<b>4,027</b>	<b>39.8 ± 5.3</b>		<b>7,751</b>	<b>248.4 ± 25.8</b>		<b>10,382</b>	<b>106.7 ± 21.5</b>
Area sampled		2,007.7 yd <sup>2</sup>			577.8 yd <sup>2</sup>			1,852.2 yd <sup>2</sup>	
Sample size		101.1 yd <sup>2</sup>			31.2 yd <sup>2</sup>			97.3 yd <sup>2</sup>	
Percent of area sampled		5.0%			5.4%			5.2%	
Length of stream sampled		120 yd			54 yd			120 yd	

Table 5. (continued). Numbers and relative abundance of naiad species found at each site sampled quantitatively in the Upper Little Black River during 1980 and 1981.

Species	River Mile 42.1			River Mile 43.5		
	Percent occurrence in quadrats	Number found	Relative abundance (%)	Percent occurrence in quadrats	Number found	Relative abundance (%)
<i>Lasmigona costata</i>	2.1	2	0.4	3.4	6	3.1
<i>Tritogonia verrucosa</i>						
<i>Amblyma p. plicata</i>						
<i>Fusconafia flava</i>	8.5	8	1.6	12.4	18	9.3
<i>Cyclonaias tuberculata</i>	17.0	20	3.9			
<i>Pleurobema coccineum</i>	48.9	94	18.4	9.0	17	8.8
<i>Elliptio dilatata</i>	45.7	188	36.7			
<i>Ptychobranchus occidentalis</i>	37.2	55	10.7	3.4	5	2.6
<i>Cyrogenia aberti</i>	6.4	7	1.4			
<i>Actinonaias' ligamentina carinata</i>						
<i>Toxolasma parvus</i>	9.6	18	3.5	3.4	5	2.6
<i>Villosa i. iris</i>	10.6	13	2.5	2.8	5	2.6
<i>Villosa l. lienosa</i>	8.5	12	2.3	3.4	7	3.6
<i>Lampsilis radiata luteola</i>	1.1	1	0.2			
<i>Lampsilis ventricosa</i>	4.3	5	1.0	2.1	3	1.6
<i>Lampsilis reeviana brevicula</i>	47.9	89	17.4	53.1	127	65.8
<i>Epioblasma florentina curtisi</i>						
<b>Naiad totals</b>	<b>80.8</b>	<b>512</b>	<b>8.6 ± 7.8 c.i.</b>	<b>67.5</b>	<b>193</b>	<b>2.1 ± 0.4 c.i.</b>
<i>Corbicula leana</i>	98.9	6,216	109.8 ± 12.9	0	0	0
<b>Area sampled</b>		<b>1,167.1 yd<sup>2</sup></b>			<b>1,842.3 yd<sup>2</sup></b>	
<b>Sample size</b>		<b>59.8 yd<sup>2</sup></b>			<b>92.2 yd<sup>2</sup></b>	
<b>Percent of area sampled</b>		<b>5.1%</b>			<b>5.0%</b>	
<b>Length of stream sampled</b>		<b>55.2 yd</b>			<b>90 yd</b>	

c.i. = 95% confidence interval.

Table 6. Water quality measurements at sites sampled in the Little Black and Castor Rivers, and Flat, Cane, and Ten-Mile creeks during 1980 and 1981.

	Little Black River						Ten-Mile														
	River Mile:	29.0	31.3	31.6	32.1	36.7	37.9	39.7	40.5	41.3	42.1	43.5	Flat Creek	Cane Creek	Ten-Mile Creek	Castor River					
Water Temperature (°C)	14	10				7	6	20	21	25	7	22	17	19	17	20	16	18			
Dissolved Oxygen (mg/l)	7.7	8.8				11.0	9.9	13.0	13.0	12.1	9.9	12.2	6.6	11.0	11.0	6.6	11.0	13.0			
pH	8.0	8.0				8.5	8.5	8.5	8.5	8.5	8.0	8.5	8.5	8.0	9.0	7.5	8.5	9.0			
Alkalinity (mg/l)	88	137				120	120	120	137	137	85	154	86	85	86	86	103	103			
Total Hardness (mg/l)	154	188				154	137	154	154	171	137	171	120	171	103	103	120	120			
Turbidity (JTU's)		1.7				1.8	1.1	1.5	5.0	2.0	1.6	4.5	8.0	5.4	4.0	2.5	2.6	1.1			
Specific Conductance (µmhos/cm <sup>2</sup> )		158				129	125	142		117	110				51						
Gradient (ft./mi.)	2.4	6.0	6.0	6.0	6.0	7.7	5.7	5.7	10.0	10.0	10.0	23.3	30.8	2.5	5.6	5.9	7.2	0.9	0.9	2.4	5.9

Table 7. Water quality measurements at six sites in the Upper Little Black River during April, 1981 through January, 1982.

	River Mile: 33.6	36.7	37.9	39.7	42.1	43.5
<u>April 10, 1981</u>						
Water temperature (°C)	20	20	20	19	18	18
Dissolved oxygen (mg/l)	9.9	7.7	7.7	7.7	7.7	7.7
pH	8.5	8.5	8.0	8.0	8.0	8.0
Alkalinity (mg/l)	86	86	86	86	68	68
Total hardness (mg/l)	137	137	120	120	103	103
Turbidity (JTU'S)	-	-	-	-	-	-
<u>June 21, 1981</u>						
Water temperature (°C)	23	24	24	23	23	23
Dissolved oxygen (mg/l)	7.7	6.6	5.5	6.6	6.6	6.6
pH	8.0	7.5	7.5	7.5	7.5	7.5
Alkalinity (mg/l)	68	68	68	51	68	51
Total hardness (mg/l)	120	120	120	103	103	86
Turbidity (JTU'S)	-	-	-	-	-	-
<u>August 8-12, 1981</u>						
Water temperature (°C)	22	23	23	24	-	26
Dissolved oxygen (mg/l)	7.7	6.6	5.5	6.6	-	7.7
pH	8.5	8.0	8.0	8.5	-	8.0
Alkalinity (mg/l)	120	120	137	137	-	120
Total hardness (mg/l)	171	205	171	154	-	154
Total nitrogen (mg/l)	0.185	0.159	0.217	0.155	0.142	0.173
Total phosphorus (mg/l)	0.034	0.036	0.030	0.034	0.033	0.030
Turbidity (JTU'S)	3.8	3.5	3.6	4.1	2.2	2.7
<u>October 19-22, 1981</u>						
Water temperature (°C)	13	14	15	13	15	15
Dissolved oxygen (mg/l)	7.7	7.7	7.7	8.8	8.8	7.7
pH	8.5	8.5	8.5	8.0	8.0	8.5
Alkalinity (mg/l)	68	68	34	68	68	51
Total hardness (mg/l)	188	171	188	171	171	154
Total nitrogen (mg/l)	0.118	0.104	0.118	0.152	0.105	0.145
Total phosphorus (mg/l)	0.018	0.008	0.016	0.020	0.002	0.021
Turbidity (JTU'S)	3.0	1.3	2.5	1.4	3.3	4.9
<u>January 12, 1982</u>						
Water temperature (°C)	0	0	-	0	0	0
Dissolved oxygen (mg/l)	12.1	12.1	-	11.0	13.2	12.1
pH	7.5	7.5	-	7.5	7.5	7.0
Alkalinity (mg/l)	0	0	-	0	0	0
Total hardness (mg/l)	120	103	-	103	103	103
Total nitrogen (mg/l)	0.470	0.510	-	0.517	0.517	0.617
Total phosphorus (mg/l)	0.011	0.014	-	0.011	0.012	0.015
Turbidity (JTU'S)	3.5	3.0	-	2.6	2.7	2.3
Conductivity	165	150	-	135	145	120

Quarterly measurements of water quality made during 1981 and early 1982 revealed little difference among the 5 sites that were quantitatively sampled in the Upper Little Black River (Table 7). Alkalinity, total hardness, and water temperature were highest during summer and lowest during winter, dissolved oxygen and total nitrogen were highest during winter, and pH was lowest during winter.

#### Flat Creek

Fifteen species of naiades were found in Flat Creek at the 2 sites sampled (Table 1). No E. f. curtisi were found. No attempt was made to estimate naiad densities in Flat Creek.

#### Castor River

Seventeen naiad species were found at the 4 sites sampled in the Castor River during 1980 and 1981 (Table 1). Although E. f. curtisi has been found as recently as 1978 in the Castor River at Zalma (river mile 9.8), none were found at that site or at the other 3 sites sampled during 1980. Habitat at river miles 20.0 and 30.3, upstream from Zalma, appeared to be too unstable for E. f. curtisi. The Castor River at river mile 3.6, downstream from Zalma, is a slow, sluggish, turbid lowland stream and not suitable habitat for E. f. curtisi.

Estimates of naiad density in the Castor River at river mile 9.8 revealed an average of 12.3 naiades and 242.2 Corbicula leana per square yard (Table 8). Nine species were found in quadrats sampled at this site. Elliptio dilatata (76.9%) and Actinonaias ligamentina carinata (9.5%) comprised 86.4% of the naiades found.

Table 8. Numbers and relative abundance of naiad species found at River Mile 9.8 in the Castor River during 1980.

Species	Percent occurrence in quadrats	Number found	No./yd <sup>2</sup>	Relative abundance(%)
Lasmigona costata	1.1	1	*	0.4
Quadrula metanevra	4.3	5	0.1	1.8
Amblema p. plicata	12.0	14	0.2	5.1
Fusconaia flava	2.2	2	*	0.7
Pleurobema coccineum	2.2	2	*	0.7
Elliptio dilatata	52.2	210	3.4	76.9
Cyprogenia aberti	9.8	11	0.2	4.0
Actinonaias ligamentina carinata	21.7	26	0.4	9.5
Ligumia recta	<u>2.2</u>	<u>2</u>	<u>*</u>	<u>0.7</u>
Naiad Totals	64.1	273	4.5	100.0
Corbicula leana	95.6	14,788	242.2	
Area sampled		2051.6 yd <sup>2</sup>		
Sample size		61.1 yd <sup>2</sup>		
Percent of area sampled		3.0%		
Length of stream sampled		102 yd		

\* = less than 0.1%

Cane Creek

Thirteen species of naiades were found at the 3 sites sampled in Cane Creek during 1980 and 1981 (Table 1). No E. f. curtisi were found. Oesch (in press) reported taking E. f. curtisi from Cane Creek at Highway M, but none were found at that site during this study. This species is very uncommon in Cane Creek and it probably occurs in only a short reach of this stream. Naiad densities were not estimated in Cane Creek.

Ten Mile Creek

Lampsilis reeviana brevicula was the only naiad species found at the single site sampled in Ten Mile Creek during 1980; no E. f. curtisi were found (Table 1).

SUMMARY AND RECOMMENDATIONS

Naiad collections were made at 11 sites in the Upper Little Black River, 2 sites in Flat Creek, 4 sites in the Castor River, 3 sites in Cane Creek, and at 1 site in Ten Mile Creek to further delineate the distribution of Epioblasma florentina curtisi in these streams. Epioblasma florentina curtisi was not found in Flat Creek or Ten Mile Creek, or at any new sites in Cane Creek or the Castor River. Epioblasma florentina curtisi was found 1.2 miles further upstream in the Upper Little Black River than previously reported, and its known range now extends from river mile 33.6 to river mile 39.7.

Epioblasma florentina curtisi was found in slow (less than 0.02 yd/sec. to 0.26 yd/sec. at bottom) current in 4 to 30 inches of water in a sand and gravel to gravel, cobble, and boulder substrate. The mean particle size of the substrate where E. f. curtisi was found, based on substrate analysis,

ranged from small gravel to cobble. The particle size and organic composition of the substrate where E. f. curtisi was found was similar to that where it was not found in the Upper Little Black River.

The 20 male and 3 female E. f. curtisi found during 1980 and 1981 ranged from 0.6 to 1.5 inches long and were determined to be from more than 4 to over 12 years old.

An average of 5.1 naiades and 74.4 Corbicula leana per square yard were found in the Upper Little Black River at the 5 sites where estimates of naiad densities were made. At 3 sites where E. f. curtisi is known to occur, an average of less than 0.1 E. f. curtisi per square yard were found.

The abundance, relative abundance, and species composition of naiades were similar at sites where E. f. curtisi occurred and sites where this species did not occur in the Upper Little Black River. Measures of water quality were similar at sites where E. f. curtisi occurred and sites where this species did not occur.

Epioblasma florentina curtisi, in both the Little Black River and other streams from which it is known, occurs in stream reaches which represent transition areas between headwater and lowland stream reaches. It is found in stream reaches of order 4 to order 7, with gradients from 0.9 to 8.0 feet per mile, in stable substrates of sand and gravel to gravel, cobble and boulder, in riffles or runs.

Habitat and population characteristics examined during this study do not explain why E. f. curtisi occurs where it does in the Upper Little Black River nor do they enable us to predict what impact proposed S.C.S. impoundments might have on this species in the Upper Little Black River. This study does underline, however, the importance of these populations to

the species as a whole. The populations of E. f. curtisi in the Upper Little Black River are the largest known populations of this species. The Upper Little Black River between river miles 33.6 and 39.7 should be considered essential habitat for the survival of E. f. curtisi unless ongoing studies (Buchanan 1982) of the distribution, abundance, and status of this species in Missouri reveal larger populations of E. f. curtisi elsewhere.

Ongoing studies on nearby Fourche Creek in southwestern Ripley County, Missouri may reveal some of the downstream changes which will occur in a small stream after impoundment. Fajen (1979) is studying the fish populations and physical characteristics at 7 sites on Fourche Creek, and the United States Geological Survey is monitoring flow and water temperature.

Future research in the Upper Little Black River should concentrate on establishing baseline water quality data for a period of several years, and further study of the life history and populations of E. f. curtisi. Populations of E. f. curtisi should continue to be monitored to determine population trends.

#### REFERENCES

- Buchanan, A. C. 1979. Mussels (Naiades) of the Little Black River Basin in Missouri and Arkansas. United States Department of Agriculture, Soil Conservation Service, Columbia, Missouri. 69 pp.
- Buchanan, A. C. 1982. Distribution, abundance, status, and habitat of the Curtis Pearly Mussel, Epioblasma florentina curtisi, in southeastern Missouri. Endangered Species Project SE-3-1. Performance Report. 5 pp.
- Fajen, O. F. 1979. An evaluation of the United States Soil Conservation Service, Fourche Creek Watershed Project. Dingell-Johnson Project F-1-R-27. Interim Report. 18 pp.
- Inman, D. L. 1952. Measures for describing the size distribution of sediments. Journal of Sedimentary Petrology. 22(3):125-145.
- Oesch, R. D. In Press. The naiades of Missouri. Missouri Department of Conservation.

An Evaluation of the Potential Impacts of SCS Structures B-9 and  
C-7 Upon the Curtis Pearly Mussel, Epioblasma florentina curtisi  
(Utterback 1915), in the Upper Little Black River, Missouri.

Alan C. Buchanan  
Missouri Department of Conservation  
1110 College Avenue  
Columbia, Missouri 65201

December, 1983

## Introduction

Epioblasma florentina curtisi is one of those species which is on the verge of extinction. Based on recent information, E. f. curtisi occurs in less than 15 miles of streams in Missouri and nowhere else in the world. Without proper management and protection of this species and its habitat, it is unlikely to survive this century. E. f. curtisi was listed as "Endangered" on the United States List of Endangered and Threatened Wildlife and Plants (United States Department of the Interior, Fish and Wildlife Service 1976) on July 1, 1975. This review is based upon information regarding the distribution and biology of E. f. curtisi, information on the characteristics of proposed structures B-9 and C-7, and literature on the impacts of impoundments on the physical, chemical and biological characteristics of streams.

Most reports documenting impacts of impoundments on aquatic organisms downstream dealt with large deep reservoirs with either surface (epilimnial) or bottom (hypolimnial) releases. Deep reservoirs with epilimnial releases generally discharge water which is warmer and lower in nutrients than waters downstream. Reservoirs with hypolimnial releases generally discharge water which is higher in nutrients and dissolved inorganic matter, lower in oxygen, and colder than waters downstream. Proposed structures B-9 and C-7 would create relatively small (298 and 136 acres, respectively), shallow (26 and 18 feet maximum depth, respectively), reservoirs with multilevel discharges (United States Department of Agriculture, Soil Conservation Service 1975). The potential impacts of these structures on the water quality of the Little Black downstream are thus expected to be much less drastic than those of large, deep reservoirs with single-level releases.

This evaluation is divided into two parts: 1) the potential impacts of structures B-9 and C-7 on the chemical, physical, and biological characteristics of the Little Black River downstream; and 2) the relative importance of Little Black populations of E. f. curtisi to the survival of the species as a whole.

Impacts of Structures B-9 and C-7 on Chemical, Physical, and Biological Characteristics of the Little Black River Downstream.

Chemical Factors

Dissolved Oxygen

Dissolved oxygen levels downstream from structures B-9 and C-7 would be at or near saturation. Although both reservoirs would stratify during summer months, water would be released from the reservoirs near the surface where dissolved oxygen levels would be expected to be high enough to support aquatic life. Turbulence immediately below the dam would restore any low-oxygen water released to saturation within a short distance downstream.

Alkalinity, Hardness, pH, and Conductivity

The impact which structures B-9 and C-7 would have on alkalinity, hardness, and pH downstream is unknown. The Little Black River is relatively well-buffered (alkalinity 34 to 137 ppm). Alkalinity dropped to 0 ppm in January, 1982 in the study reach because of rainfall runoff at a time when the ground was frozen. During normal flow conditions, however, the Little Black probably remains well-buffered. Little Black waters are also moderately hard (86 to 205 ppm total hardness) with a pH usually above 8.0 (7.0 to 8.5).

The reservoirs created by both structures B-9 and C-7 would be expected to stratify during summer. During stratification in eutrophic hard water reservoirs (which both of these reservoirs would be) there usually is a reduction of total carbon dioxide in the epilimnion due to photosynthesis by vascular plants and algae (Hutchinson 1957; Ruttner 1963; Wetzel 1975). With a removal of carbon dioxide there is a precipitation of calcium carbonate and a concurrent rise in pH. The pH would not be expected to greatly exceed 9.0, however. In the hypolimnion there is a progressive increase in carbon dioxide, due to respiration and release from the sediments, with a resultant reduction in pH and rise in bicarbonate. Again, the pH would be expected to remain relatively moderate and not drop below 7.0.

Since water would be drawn from the epilimnion in both reservoirs and released downstream, this water would be near or above the pH of water in the stream below. Total hardness and alkalinity in the water released from this level would be less than downstream. The conductivity of the water released from structures B-9 and C-7 would also be expected to be lower than ambient levels downstream since reservoirs generally reduce the conductivity of water. The effect that an increase in pH and a reduction in alkalinity, hardness, and conductivity would have on benthic macroinvertebrates in Little Black River is expected to be minimal.

There would also be fluctuation in levels of pH, alkalinity, hardness, and conductivity in the reservoirs during the seasons. During spring and fall overturn, when there is a mix of surface and bottom water within the reservoirs, levels of pH, alkalinity, hardness, and conductivity would be similar from surface to bottom. Thus, in water released from the reservoirs, the pH would be lower and the alkalinity, hardness, and conductivity higher

than during times of stratification. The effect that these annual fluctuations would have on the biota of the Little Black River 3.5 to 6 miles downstream from these structures is not known.

Below Fourche Dam #1 on Fourche Creek, a tributary to Current River in Ripley County, Missouri, there was a reduction in conductivity, total alkalinity, and total hardness within and immediately below the lake as compared to above and 5 miles below the lake (Missouri Department of Conservation memo from Rich Wehnes to Stan Michaelson dated November 28, 1978). There was no change in pH within the reservoir. This information is based on a single set of measurements made in November, 1978.

#### Nutrients

Reservoirs generally serve as nutrient sinks, with nutrients in the incoming waters being trapped in the bottom sediments. A review of the literature suggests that there would be a reduction of nutrients in waters released from structures B-9 and C-7 in comparison to levels in the Little Black River entering the reservoirs. Levels of nitrogen, phosphorus, iron, and zinc would probably be reduced.

Turner et al. (1983) reported a mean loss over a 3-year period of 44.8% of the total phosphorus and 10% of the total nitrogen from water entering Lake Talquin in northwestern Florida. The amount of phosphorus and nitrogen lost to the reservoir was inversely related to the flushing rate in any given year. The flushing rate in Lake Talquin ranged from 8 to 16 times per year. Lake Talquin had no significant effect on silicates or chloride in water passing through it.

In a review of the literature, Turner et al. (1983) found that phosphorus and nitrogen are frequently reduced by reservoirs. They found that phosphorus

was reduced by 25% or more from the inflow to the outflow of 20 of 27 reservoirs examined. In 3 of the 27 reservoirs phosphorus was reduced by more than 50% and in 2 others by more than 75%. Nitrogen was also retained by reservoirs but to a lesser extent than phosphorus. Ten of thirteen reservoirs examined by Turner et al. (1983), retained 10% or more of the total nitrogen entering them. Seven reservoirs retained 25% or more of the total nitrogen entering them. Although there was a trend toward greater retention of nitrogen and phosphorus in reservoirs with longer retention times, especially in those with retention times of more than a year, there was no significant correlation between retention time and retention of nitrogen and phosphorus by reservoirs.

Several other studies report significant retention of nitrogen and phosphorus by reservoirs. In Bighorn Lake, Montana (Soltero et al. 1973, 1974) there was a reduction of phosphorus by 86% and of nitrogen by 25% from the inflow to the outflow. Higgins (1978) reported a reduction of total nitrogen and total phosphorus from the inflow to the outflow in five Tennessee Valley Reservoirs over a 10-year period. Although nutrients are frequently reduced by reservoirs, they are not necessarily reduced to levels that would be limiting to the streams below.

Frequently the reduction of nutrients in the outflow of water from a reservoir is seasonal, generally during summer stratification. During fall overturn, there is often an increase in phosphorus and nitrogen in the outfall due to the mixing of hypolimnial water, which is higher in these nutrients, with water from the epilimnion. Thus there are seasonal fluxes of nutrients to the stream below the reservoirs.

The impact of B-9 and C-7 upon the nutrient levels of the Little Black River 3.5 to 6 miles downstream is unknown. Due to the expected short

retention time of water in these reservoirs, there will probably be a minimal retention of nutrients. The seasonal fluctuation of nutrients in water released from the reservoirs would affect benthic macroinvertebrates immediately below the dams, but might have little effect on benthos 3.5 to 6 miles downstream.

#### Dissolved Solids

Higgins (1978) reported that total dissolved solids were lower in the outflow than in the inflow of five Tennessee Valley Authority reservoirs. Total dissolved solids were slightly lower in the outflow than the inflow in Bighorn Lake, Montana (Soltero et al. 1973, 1974). Dissolved solids will be reduced by the reservoirs formed by structures B-9 and C-7. The impact, if any, downstream would be minimal.

#### Physical Factors

##### Temperature

Due to the arrangement of release ports on structures B-9 and C-7, little or no change in downstream temperature would be expected. The only direct effects may be a retarding of spring warming and fall cooling. Retarding the spring increase in water temperatures might impact E. f. curtisi if it occurs during the spawning period for this species. The likelihood of impact is proportional to the amount spring warming is retarded. Studies on other small reservoirs, such as those on Fourche Creek may provide additional information on the impact of small reservoirs on the rate of spring warming and fall cooling of receiving streams. Both reservoirs would be expected to stratify thermally in the summer and possibly in the winter.




Figure 1. Little Black River Basin, Missouri.

Summer surface temperatures in the reservoirs would be significantly higher than that of inflowing waters. Consequently, the reservoirs would support a different biotic fauna than the stream.

#### Downstream Flows

Peak flows would be reduced by construction of structures B-9 and C-7. Peak flows 2 1/2 miles downstream from Highway K, (cross section 19, Fig. 1) near the reach where E. f. curtisi occurs, would be reduced from 12,500 cfs to 2,800 cfs for a 25-year frequency flood, and from 3,000 cfs to 1,400 cfs for a 1-year frequency flood (United States Department of Agriculture, Soil Conservation Service, unpublished data). Bank full at this site is 3,000 cfs. Two miles further downstream (cross section 21, Fig. 1), in the reach where the best populations of E. f. curtisi occur, peak flows would be reduced from 14,000 cfs to 3,600 cfs for a 25-year frequency flood, and from 1,325 cfs to 900 cfs for a 1-year frequency flood. Bank full at this latter site is 1,830 cfs.

Although peak flows would be reduced by structures B-9 and C-7, duration of flood flows would be increased significantly. The project is designed to reduce the volume and duration of out-of-bank flows, but would increase the length of time of bank-full and above normal in-bank flows. The change in duration of a 25-year flood event which would result from the projects is as follows (cross section 17, Fig. 1): Without structures B-9 and C-7 it takes 10.25 hours for the waters to overflow the banks. Flows remain out-of-bank for 19.75 hours, and return to normal (7 cfs) 43.5 hours after the initial increase. With structures B-9 and C-7 in place, it would take 13.5 hours for the waters to overflow the banks. They would remain out-of-bank for 10

hours, and return to 450 cfs in 72.5 hours. It could take up to 10 days for flows to return to normal (7 cfs) after the beginning of the flood event (United States Department of Agriculture, Soil Conservation Service unpublished data). The change in duration of flows for 10- and 100-year floods would be very similar to that described for the 25-year flood event.

The impact of a combination of reduced peak flows and reduced duration of out-of-bank flows, and an increase in duration of above normal in-bank flows is unknown. While a reduction in peak flows would tend to reduce scouring downstream, increased duration of above normal in-bank flows would have the opposite effect. Based on unpublished data provided by the United States Department of Agriculture, Soil Conservation Service, bedload transport during 25-year and 1-year flood events would be reduced by factors of 6 and 3, respectively, at cross section 19 (Fig. 1). This would result in aggradation of the streambed below the proposed reservoirs if a source of bed material were available. Structures B-9 and C-7 and other structures in the basin will severely reduce the source of bedload material, however, and degradation is expected.

While a reduction in peak flows would reduce scouring of the streambed and banks and reduce exchange of nutrients and organic matter between the stream and floodplain, extended duration above normal in-bank flows would tend to increase bed scouring and bank scouring and sloughing. The net impact upon instream vegetation and bank stability is not known. With a decrease in out-of-bank flows, riparian vegetation will probably change to species characteristic of more advanced successional stages.

The net impact upon benthic macroinvertebrates downstream is also not known. Evidence from at least one previous study indicates that changes in flows, particularly elimination of seasonal peak flows may cause significant changes in the species composition and abundance of benthic macroinvertebrates.

Williams and Winget (1979) found an increase in aquatic macrophytes and epiphytic algae below Soldier Creek Dam, a small impoundment on the Strawberry River, a high gradient stream in Utah. Although there was no change in diversity as indicated by the Shannon Weaver Index, there was a change in taxonomic composition of benthic macroinvertebrates. Numbers of low-flow tolerant species increased significantly, while some species disappeared entirely. They attributed these changes to a reduction in habitat heterogeneity due to reduced seasonal flows. A reduction in peak flows in the Strawberry River also allowed the proliferation of beaver dams which were previously washed out annually by spring flooding. Beaver dams backed water over riffles, reducing riffle flow velocity and allowing the accumulation of finer sediments in riffles.

An increase in duration of above normal in-bank flows during a flood event would result in an increase in duration of above normal turbidity, which may limit the spawning success of E. f. curtisi. This species spawns in the spring, when high flows are most common. Extended periods of valve closure by naiades because of increased turbidity may reduce the time available to release glochidia (larval naiades). Glochidia and juvenile naiades are more susceptible to silt and other turbidity than adults and would be adversely affected by extended periods of above normal levels of turbidity.

#### Suspended Sediment

Water released downstream from structures B-9 and C-7 would contain less sediment than the water presently contains. A reduction in sediment downstream could, ultimately, alter the composition of the substrate. Water released from the above structures would be "sediment hungry." This could increase bed-cutting and reduce the fines in the substrate downstream. Either

change may adversely impact naiades because substrate quality appears to be critical for most species. The mean particle size of the substrate where E. f. curtisi occurs ranges from small gravel to pebble (0.04 to 1.25 inches in diameter).

### Biological Changes

The most important, yet most difficult to predict, downstream changes which may occur in a stream after construction of an impoundment are changes in the biological constituent of the stream. Potential changes include increases or reduction in numbers of organisms present, a change in species dominance, and addition or loss of one or more species.

An impoundment creates an environment suited to lake species but poorly suited for many riverine species which occur in the stream before impoundment. Thus, impoundments contain a different array of organisms than streams. Consequently, the organisms which pass from a reservoir into the stream below are generally much different than those occupying the stream.

Cummins (1979), in a discussion of stream ecosystems as continuums, discussed impacts which impoundments have upon stream reaches downstream from the impoundments. A stream is a continuum from its headwaters to its mouth. In its headwaters (orders 1 to 3) the stream is most impacted by inputs from outside the channel. As you move downstream, the influx of materials from upstream in-channel sources becomes progressively more important and out-of channel inputs progressively less important. While coarse particulate matter processing is most important in the upper reaches of a stream, fine particulate organic matter processing is most important in large rivers. Impounding a stream has the effect downstream similar to

increasing the stream order; the continuum is interrupted. Organisms and organic matter processing below an impoundment change to resemble that of communities much farther downstream. Impoundments impart a stability to downstream reaches which is more characteristic of larger stream reaches. This may alter not only the plankton and benthos of the downstream reach, but also the fish which feed upon them. E. f. curtisi has been found only in stream reaches which serve as transition areas between headwater and lowland reaches. Therefore, any change in stream characteristics which resemble a change in stream order would impact this species.

Several authors have described differences between the organisms present immediately downstream from a reservoir and those in unimpacted portions of the stream further downstream. Lowe (1979) found that phytobenthos downstream from reservoirs increased because of increased light penetration and regular flows. Armitage and Capper (1976) documented the transport of lake zooplankton downstream from reservoirs. They found that the amount of lake plankton in a stream below a reservoir was inversely related to the distance below the reservoir, turbulence, and amount of vegetation present. The relative size of the reservoir and the stream into which it emptied, and the season of the year were also important. In nine cases discussed, lake species comprised from 100% of the plankton in Maple Creek 35 feet downstream from Douglas Lake to 1% in the Huron River 4.5 miles downstream from Portage Lake. In the Marais River, lake species comprised 96% of the zooplankton 7.75 miles downstream from Tiber Reservoir.

The type of plankton and other microorganisms present in a stream affects not only the food available for naiades, which feed on plankton and organic matter, but also the kinds of fish present which act as potential hosts for

the glochidia of naiades. Changing the amounts and types of microorganisms may thus impact the survival of naiades downstream. The reach of the Little Black River which supports E. f. curtisi is approximately 3.5 and 6.0 miles, respectively, downstream from the proposed sites of B-9 and C-7. It is not known with certainty whether these projects would impact the biological component of the Little Black this far downstream. Unfortunately, neither the host(s) nor the specific food of the Curtis Pearly Mussel are known.

Epioblasma, the genus of which the Curtis Pearly Mussel is a member, is the most recently evolved genus of freshwater naiades, and one of the most specialized. Species of this genus are found only in riffles, in stable substrates, in waters of relatively high quality. Because members of this genus are so specialized, they have been severely impacted by human activities since the Industrial Revolution. Only one species in the genus, Epioblasma triquetra, is not in immediate danger of extinction. Holden (1979) in reference to fish in regulated streams, stated that the most specialized species of fish disappear first when a stream is altered. This also appears to be true in the case of naiades. If structures B-9 and C-7 produce any significant downstream changes, E. f. curtisi would be one of the first species impacted.

#### The Relative Importance of the Little Black Populations of Epioblasma florentina curtisi

---

Any impacts which structures B-9 and C-7 might have on populations of Epioblasma florentina curtisi in the Upper Little Black River become more significant when the relative importance of the Little Black River population of E. f. curtisi to the species as a whole is considered.

E. f. curtisi has not been found outside of Missouri in recent times. The best populations known of this species occur in approximately 6 miles of the Upper Little Black River. During the past 5 years, over 60 specimens of E. f. curtisi have been examined in the Little Black River, more than have been found previously throughout its entire range. Outside of the Little Black River, specimens of this species have been found at only one site each in Cane Creek (Oesch in press) and the Castor River, and at several sites in Black River (Dr. David H. Stansbery, Ohio State University Museum of Zoology personal communication) during the past 15 years.

A recent survey of E. f. curtisi in southern Missouri (Buchanan 1983) failed to expand the range of this species. More importantly, no fresh material of this species was found during extensive sampling in Black River and E. f. curtisi probably no longer occurs in Black River. The last fresh material of E. f. curtisi collected from Black River was collected in 1970 (Dr. David H. Stansbery personal communication). E. f. curtisi is now known to occur only in 6.1 miles of the Upper Little Black River, 5 miles or less of the Castor River, and one site in Cane Creek. Population levels remain very low in Castor River and Cane Creek.

The disappearance of E. f. curtisi from Black River may be tied to human activities, including the construction of Clearwater Dam, which was closed in 1948. Since closure of Clearwater Dam, flow characteristics in Black River downstream have changed drastically. Hydrographic records at Leeper, 6.0 miles downstream from Clearwater Dam, reflect these changes (United States Geological Survey unpublished data). Between 1922 and 1948, the maximum annual flow was less than 10,000 cubic feet per second (cfs) only 8 times and was less than 5,000 cfs only twice. Since construction of Clearwater

Dam, the maximum annual flow at Leeper has never exceeded 10,000 cfs and has exceeded 5,000 cfs only twice. Periodicity of flows has been altered also. Mean daily flows at Leeper during April, May, and June were reduced by 357, 124, and 335 cfs, respectively, by Clearwater Dam. Mean daily flows during July, August, and September have been increased by 198, 160, and 126 cfs, respectively, by the dam. Clearwater Dam has also resulted in extremely low flows by Leeper. Between 1922 and 1948, annual minimum flows dropped below 200 cfs only 7 times, and the lowest flow was 133 cfs in August, 1934. Between 1949 and 1979 annual minimum flows dropped below 200 cfs 14 times and below 100 cfs twice. The lowest flows recorded were 70 cfs in November, 1959 and 62 cfs in September, 1966. The above flow changes could impact the time and bottom area available for fish, potential hosts for E. f. curtisi and other naiades, to spawn. Very low flows may expose concentrations of naiades, causing desiccation, temperature stress, and increased susceptibility to predators. Changes in flow periodicity may also impact reproduction in benthic macroinvertebrates.

#### Summary and Recommendations

1. Structures B-9 and C-7 are not expected to have a significant impact upon water temperature or the levels of dissolved oxygen downstream.
2. The impact of B-9 and C-7 upon nutrient levels downstream would depend upon annual temperature and rainfall regimes, and characteristics of the reservoirs created, including flushing rates, level and duration of stratification, and other factors. Nutrient levels in the outflow from these reservoirs would change during each year as the reservoirs stratify and destratify from season to season.

3. Although stratification related changes in pH and alkalinity within the reservoirs created by B-9 and C-7 would be expected, the Little Black is relatively well-buffered and no significant impacts are expected downstream.
4. Structures B-9 and C-7 would act as sediment traps, reducing the load of sediments passing downstream. This may have an impact on substrate composition and stability downstream. The extent of downstream impact, if any, is not known.
5. The impact of the combination of reduced peak flows and extended duration of above normal in-bank flows created by structures B-9 and C-7 is not known. While reducing peak flows and the duration of out-of-bank flows would tend to reduce scouring and bank-cutting, extending the duration of above normal in-bank flows would have the opposite effect. The net impact would probably be degradation of the streambed downstream due to entrapment of sediments behind B-9 and C-7 and other structures in the basin, possibly altering the composition of the substrate.
6. Longer duration above normal in-bank flows would be expected to be accompanied by longer periods of above normal turbidity. Since peak flows in the Little Black River normally occur in the spring when E. f. curtisi is releasing glochidia, extended duration above-normal flows may result in longer periods of above-normal turbidity at a very crucial time for this species. Glochidia and juvenile naiades are particularly sensitive to silt, and higher turbidity may reduce the chances for successful reproduction.
7. Impoundments frequently alter the biota of the streams into which they empty. Previous studies have documented changes in periphyton, vegetation, plankton, benthic macroinvertebrates, and fish populations

downstream from reservoirs. Water passing from reservoirs usually contains organisms quite different from those occupying flowing waters downstream. Changes in the plankton available might alter the food supply and a change in fish populations might reduce the number of potential hosts for E. f. curtisi.

8. Populations of E. f. curtisi in the Upper Little Black River are the best known. This species has experienced a serious reduction in numbers and range since the early 1900s, primarily due to dam construction. Outside of the Little Black River, E. f. curtisi occurs in very low numbers in Castor River and Cane Creek, and nowhere else. Disappearance from Black River since 1970 may be at least partially attributed to altered flows below Clearwater Dam since its construction in 1948.
9. Sensitive species such as E. f. curtisi are generally the first species to disappear when a change takes place in a stream. Any change which impacts E. f. curtisi in the Upper Little Black River will impact the existence of this species as a whole. Disappearance of this species from the Little Black River will probably lead to its extinction.
10. Since construction of structures B-9 and C-7 on the Upper Little Black River may impact E. f. curtisi, they should not be built until the survival of this species can be assured.

References Cited

- Armitage, P. D. and M. H. Capper. 1976. The numbers, biomass and Hydra from Cow Green Reservoir (Upper Teesdale). *Freshwater Biology* 6:425-432.
- Buchanan, A. C. 1982. A study of Epioblasma florentina curtisi (Utterback 1915), the Curtis Pearly Mussel in the Upper Little Black River, Missouri, U. S. Department of Agriculture, Soil Conservation Service. 11 pp.
- Buchanan, A. C. 1983. A survey of the Curtis Pearly Mussel, Epioblasma florentina curtisi (Utterback 1915), in southern Missouri. Missouri Department of Conservation. Draft Report.
- Cummins, K. W. 1979. The natural stream ecosystem. Pages 7-24 in J. V. Ward and J. A. Stanford, editors. *The Ecology of Regulated Streams*. Plenum Press.
- Higgins, J. M. 1978. Water quality progress in the Holston River Basin. Water Quality and Ecology Branch, Division of Environmental Planning. Tennessee Valley Authority, Chattanooga, Tennessee. 43 pp.
- Hutchinson, G. E. 1957. *A Treatise on Limnology*. Volume 1.
- Lowe, R. L. 1979. Phytobenthic ecology and regulated streams. Pages 25-35 in J. V. Ward and J. A. Stanford, editors. *The Ecology of Regulated Streams*. Plenum Press.
- Oesch, R. D. In Press. The Naides of Missouri. Missouri Department of Conservation.
- Ruttner, F. 1963. *Fundamentals of Limnology*. Third edition, University of Toronto Press, Toronto, Canada.
- Soltero, R. A., J. C. Wright, and A. A. Horpestad. 1973. Effects of impoundment on the water quality of the Bighorn River. *Water Resources* 7:343-354.

Turner, R. R., E. A. Laws, and R. C. Harriss. 1983. Nutrient retention and transformation in relation to hydraulic flushing rate in a small impoundment. *Freshwater Biology* 12:113-127.

United States Department of Agriculture, Soil Conservation Service. 1975. Upper Little Black Watershed final work plan for watershed protection, flood prevention, and recreational development. 100 pp.

Williams, R. D. and R. N. Winget. 1979. Macroinvertebrate response to flow manipulation in the Strawberry River, Utah (U.S.A.). Pages 365-376 in J. V. Ward and J. A. Stanford, editors. *The Ecology of Regulated Streams*.

Wetzel, R. G. 1975. *Limnology*. W. G. Saunders Company, Philadelphia, Pennsylvania. 743 pp.

APPENDIX C  
LIST OF REVIEWERS

Technical draft sent for review to:

Ronald D. Oesch  
9 Hill Drive  
Glendale, MO 63122

Dr. Leslie Holland  
National Fishery Research Laboratory  
P. O. Box 818  
LaCrosse, WI 54601

Dr. Richard Neves  
Virginia Cooperative Fisheries Research Unit  
Virginia Polytechnical Institute  
106 Cheatham Hall  
Blacksburg, VA 24061

Steve Ahlstedt  
Tennessee Valley Authority  
Office of Natural Resources  
Norris, TN 37828

Dr. David Stansbery  
Ohio State University Museum of Zoology  
Ohio State University  
Columbus, OH 43210

Dr. Andrew Miller  
Environmental Laboratory  
U.S. Army Engineers Waterways Experiment  
Station  
P. O. Box 631  
Vicksburg, MS 39180-0631

Dr. Richard Sparks  
River Research Laboratory  
Illinois Natural History Survey  
Havana, IL 62644

Dr. John C. Williams  
Dept. of Biological Sciences  
Eastern Kentucky University  
Richmond, KY 40475

Dr. Edward M. Stern  
Department of Biology  
University of Wisconsin  
Stevens Point, WI 54481

Dr. Marc Imlay  
National Fishery Research Laboratory  
Rt. # 1  
Columbia, MO 65201

Mr. Sam Fuller  
1053 Mapleton Avenue  
Suffield, CN 06078

Paul W. Parmalee  
Department of Anthropology  
University of Tennessee  
Knoxville, Tennessee 37916

Ray Matheny  
Hazard Evaluation Division  
(OPP)-TS769  
Environmental Protection Agency  
401 M Street SW  
Washington, D.C. 20460

Agency draft sent for review to:

Mr. Larry R. Gale, Director  
Department of Conservation  
Attention: Dr. James H. Wilson  
Endangered Species Coord.  
P.O. Box 180  
Jefferson City, MO 65102

Colonel Daniel M. Wilson, Dist. Engr.  
U.S. Army Corps of Engineers  
210 Tucker Boulevard North  
St. Louis, MO 63101

Mr. Larry Henson, Regional Forester  
U.S. Forest Service  
310 W. Wisconsin Avenue  
Milwaukee, WI 53203

Steve N. Wilson, Director  
Game and Fish Commission  
#2 Natural Resources Drive  
Little Rock, AK 72205

John W. Humke, Director  
Midwest Regional Office  
The Nature Conservancy  
328 E. Hennepin Avenue  
Minneapolis, MN 55414

Tennessee Valley Authority  
Mr. Bobby G. Grinstead  
Supervisor of Biological Resources  
524 Union Avenue  
Knoxville, TN 37902

Office of Endangered Species  
U. S. Fish & Wildlife Service  
1000 N. Glebe Road  
Arlington, VA 22201

